

The exact moments of a ratio of quadratic forms in normal variables

Jan R. Magnus *

ABSTRACT. — The exact moments of $x'Ax/x'Bx$ are obtained, where x is a normally distributed vector with some mean (possibly nonzero) and positive definite covariance matrix, A is symmetric and B positive semidefinite. These moments appear as simple integrals which can be evaluated numerically in a straightforward manner. In addition, the precise conditions for the existence of the moments are found. Some related results are also reported.

Les moments exacts d'un rapport de formes quadratiques de variables normales

RÉSUMÉ. — On obtient les moments exacts de $x'Ax/x'Bx$ où x est un vecteur normal de moyenne quelconque de matrice de covariance définie positive, A est symétrique et B semi-définie positive. Ces moments sont des intégrales simples qui peuvent être évaluées numériquement de façon simple. De plus des conditions précises pour l'existence de ces moments sont fournies. D'autres résultats sont également présentés.

* J. R. MAGNUS: London School of Economics, Department of Economics, Houghton Street, Aldwych, London WC2A 2AE, Royaume Uni. My interest in the subject matter of this paper was stimulated by two recent workshops at the LSE, one by Jean-Marie DUFOUR, the other by Asraul HOQUE. I am indebted to both. Critical comments by Steve ALPERN, Risto HEIJMANS and Roald RAMER are gratefully acknowledged. A preliminary version of this paper was presented at the ESRC Econometric Study Group Conference in Bristol, July 1986. I am very grateful for the constructive comments by the participants of this conference, in particular Bill FAREBROTHER, John NANKERVIS, Peter PHILLIPS and Gene SAVIN. I also thank the two anonymous referees for their constructive remarks.

1 Introduction

Econometric or time-series estimators often take the form of ratios of quadratic forms. A very simple example is provided by the first-order autoregressive process

$$y_t = \beta y_{t-1} + u_t \quad (t=2, \dots, T)$$

where $|\beta| < 1$ and the u_t are i. i. d. $N(0, \sigma^2)$. Given a series of T observations, the least-squares estimator of β is

$$\hat{\beta} = \left[\sum_{t=2}^T y_t y_{t-1} \right] / \left[\sum_{t=2}^T y_{t-1}^2 \right]$$

which is a ratio of two quadratic forms in $y = (y_1, \dots, y_T)'$.

The purpose of this paper is to find the exact moments of such estimators, that is to say, estimators that can be written

$$x'Ax/x'Bx$$

where A is symmetric, B positive semidefinite and x is normally distributed with some mean (possibly nonzero) and positive definite covariance matrix. The procedure to obtain such moments was, in principle, given by SAWA [1978], but the evaluation of the moments is complicated by the fact that s -fold differentiation under the integral sign has to be performed in order to obtain the s th moment. As a result of this difficulty we know of only a couple of papers (DEGOOYER [1980], ALI [1984]) where third and fourth moments have been considered, and then only in the central case with zero mean; the evaluation of fifth and higher moments has never been attempted to our knowledge. In the present paper this problem is solved, thus opening the way to evaluate the exact moments in the central and the non-central case to an arbitrary order.

The literature on quadratic forms is vast; a good survey can be found in Chapters 28 and 29 of JOHNSON and KOTZ [1969-1970]. Regarding the expectation of products of quadratic forms we mention papers by LANCASTER [1954], KUMAR [1973], SRIVASTAVA and TIWARI [1976], MAGNUS [1978, 1979] and DON [1979]. Regarding the *distribution* (as opposed to the moments) of quadratic forms and ratios of quadratic forms we mention JOHN GURLAND's influential papers [1948, 1953, 1955, 1956-1957], IMHOF's [1961] important contribution, and papers by TANEJA [1976], GIDEON and GURLAND [1976], KHURI and GOOD [1977] and WALKER [1979]. For the numerical aspects the reader should consult LUGANNINI and RICE [1984]. Special cases of the *moments* of ratios of quadratic forms were recently discussed by DWIVEDI and CHAUBEY [1981] and CHAUBEY and NUR ENAYET TALUKDER [1983]. MORAN [1948] and DUFOUR and ROY [1985] gave closed-form expressions for the first and second moments of sample autocorrelations under randomness (besides results for non-normal distributions).

Finally, econometricians have been typically interested in the moments of (least-squares) estimators (SAWA [1978], HOQUE and PETERS [1986]), the distribution of forecasts arising from a first-order autoregression (PHILLIPS [1979]), and the moments of forecasts in distributed lag models (HOQUE [1985]).

The main result of the paper is stated and proved in section 5. In sections 2-4 we give five lemmas, of independent interest, which are used in the proof of the theorem in section 5. Section 6 deals with the existence of the moments and section 7 contains some remarks on computational aspects and concludes the paper.

2 Moments, Cumulants and Quadratic Forms

If x is a random variable whose s th moment $\alpha_s = E x^s$ exists, then its characteristic function $\varphi(t)$ can be expanded in a neighbourhood of $t=0$ as follows:

$$(1) \quad \varphi(t) = 1 + \sum_{h=1}^s \frac{(it)^h}{h!} \alpha_h + o(t^s)$$

where

$$\lim_{t \rightarrow 0} \frac{o(t^s)}{t^s} = 0.$$

If $\varphi(t)$ can be expanded as in (1), then $\log \varphi(t)$ may be expanded as

$$(2) \quad \log \varphi(t) = \sum_{h=1}^s \frac{(it)^h}{h!} \kappa_h + o(t^s).$$

The quantities κ_h are called the *cumulants* of the distribution of x . The following lemma, which is a simplified version of a result given by KENDALL and STUART ([1977], section 3.14, p. 70), shows that α_s is a polynomial in the cumulants $\kappa_1, \dots, \kappa_s$, and gives the coefficients of the polynomial.

LEMMA 1: If x is a random variable whose s th moment α_s exists, then its first s cumulants $\kappa_1, \dots, \kappa_s$ exist also, and α_s can be expressed as

$$(3) \quad \alpha_s = \sum_{\mathbf{v}} \frac{s!}{n_1! n_2! \dots n_s!} \left[\frac{\kappa_1}{1!} \right]^{n_1} \left[\frac{\kappa_2}{2!} \right]^{n_2} \dots \left[\frac{\kappa_s}{s!} \right]^{n_s}$$

where the summation is over all $1 \times s$ vectors $\mathbf{v} = (n_1, n_2, \dots, n_s)$ whose components n_j are nonnegative integers satisfying

$$(4) \quad n_1 + 2n_2 + \dots + sn_s = s.$$

Proof: Taking exponentials on both sides of (2) gives

$$\begin{aligned}
 \varphi(t) &= \exp \left[\sum_{r=1}^s \frac{(it)^r}{r!} \kappa_r \right] \exp(o(t^s)) \\
 &= \exp(o(t^s)) \prod_{r=1}^s \exp \left[\frac{(it)^r}{r!} \kappa_r \right] \\
 &= (1 + o(t^s)) \prod_{r=1}^s \left\{ \sum_{j=0}^s \left[\frac{(it)^{rj} (\kappa_r/r!)^j}{j!} \right] + o(t^s) \right\} \\
 &= \prod_{r=1}^s \sum_{j=0}^s \left[\frac{(\kappa_r/r!)^j}{j!} (it)^{rj} \right] + o(t^s).
 \end{aligned}$$

If, for $r=1, 2, \dots, s$, we now pick out the n_r -th term in the r -th sum such that, when multiplied together, these terms give a power of t^s , we find, upon comparison with (1),

$$\frac{\alpha_s}{s!} = \sum_{\mathbf{v}} \frac{(\kappa_1/1!)^{n_1}}{n_1!} \cdot \frac{(\kappa_2/2!)^{n_2}}{n_2!} \cdot \dots \cdot \frac{(\kappa_s/s!)^{n_s}}{n_s!}$$

and the result follows. □

In order to implement Lemma 1 for given value of s we must find the set of $1 \times s$ vectors (n_1, n_2, \dots, n_s) satisfying (4). These are easily calculated. For $s=1, 2, 3, 4$ we have

$$s=1: n_1 = 1$$

$$s=2: (n_1, n_2) = (2, 0), (0, 1)$$

$$s=3: (n_1, n_2, n_3) = (3, 0, 0), (1, 1, 0), (0, 0, 1)$$

$$s=4: (n_1, n_2, n_3, n_4) = (4, 0, 0, 0), (2, 1, 0, 0), (1, 0, 1, 0), (0, 2, 0, 0), (0, 0, 0, 1)$$

and hence the first four moments in terms of the cumulants are

$$\alpha_1 = \kappa_1$$

$$\alpha_2 = \kappa_1^2 + \kappa_2$$

$$\alpha_3 = \kappa_1^3 + 3 \kappa_1 \kappa_2 + \kappa_3$$

$$\alpha_4 = \kappa_1^4 + 6 \kappa_1^2 \kappa_2 + 4 \kappa_1 \kappa_3 + 3 \kappa_2^2 + \kappa_4.$$

For computational (and theoretical) purposes it is useful to know for given s how many vectors (n_1, n_2, \dots, n_s) there are satisfying (4). We shall address this problem in some more detail in section 7.

Our next task is to calculate the cumulants of the distribution of a quadratic form in normal variables.

LEMMA 2: Let x be a normally distributed $n \times 1$ vector with mean μ and positive definite covariance matrix $\Omega = LL'$, and let A be a symmetric $n \times n$ matrix. Then the cumulants $\kappa_1, \kappa_2, \dots$ of the distribution of $x'Ax$ are

$$(5) \quad \kappa_j = 2^{j-1} (j-1)! \{ \text{tr}(L'AL)^j + j\mu' L'^{-1} (L'AL)^j L^{-1} \mu \}.$$

COMMENT: In practice the $n \times n$ matrix L will usually be taken as (lower) triangular. But Lemma 2 (and all subsequent results) hold for any $n \times n$ matrix L satisfying $LL' = \Omega$.

Proof: Let P be an orthogonal $n \times n$ matrix and Λ a diagonal $n \times n$ matrix such that

$$P' L' A L P = \Lambda, \quad P' P = I_n.$$

Then we can write

$$x = LP(u + \omega)$$

where $u \simeq N(0, I_n)$ and $\omega = P' L^{-1} \mu$. Hence

$$x'Ax = (u + \omega)' \Lambda (u + \omega)$$

and the j -th cumulant, as given e. g. by JOHNSON and KOTZ ([1970], Chapter 29, section 3), is

$$\begin{aligned} \kappa_j &= 2^{j-1} (j-1)! (\text{tr} \Lambda^j + j\omega' \Lambda^j \omega) \\ &= 2^{j-1} (j-1)! \{ \text{tr}(L'AL)^j + j\mu' L'^{-1} P(P'L'ALP)^j P' L^{-1} \mu \} \\ &= 2^{j-1} (j-1)! \{ \text{tr}(L'AL)^j + j\mu' L'^{-1} (L'AL)^j L^{-1} \mu \}. \quad \square \end{aligned}$$

Combining Lemma 1 and Lemma 2, we obtain the moments of a quadratic form in normal variables in a readily usable form.¹

LEMMA 3: Let x be a normally distributed $n \times 1$ vector with mean μ and positive definite covariance matrix $\Omega = LL'$, and let A be a symmetric $n \times n$ matrix. Then, for $s = 1, 2, \dots$

$$(6) \quad E(x'Ax)^s = \sum_v \gamma_s(v) \prod_{j=1}^s \{ \text{tr}(L'AL)^j + j\mu' L'^{-1} (L'AL)^j L^{-1} \mu \}^{n_j}$$

where the summation is over all $1 \times s$ vectors $v = (n_1, n_2, \dots, n_s)$ whose components n_j are nonnegative integers satisfying $\sum_{j=1}^s jn_j = s$, and

$$(7) \quad \gamma_s(v) = s! 2^s \prod_{j=1}^s [n_j! (2j)^{n_j}]^{-1}.$$

1. Lemma 3 can also be obtained using zonal polynomials, see JAMES [1964], DAVIS [1979] and HILLIER and SMITH [1983].

Proof: Let $\kappa_1, \kappa_2, \dots$ be the cumulants of the distribution of $x'Ax$, and let

$$\beta_j = \text{tr}(L'AL)^j + j\mu'L'^{-1}(L'AL)^jL^{-1}\mu.$$

Then, from Lemma 2

$$(\kappa_j/j!)^{n_j} = 2^{jn_j} \beta_j^{n_j} / (2j)^{n_j}$$

and hence from Lemma 1

$$\alpha_s = s! \sum_{\mathbf{v}} \prod_{j=1}^s [n_j! (2j)^{n_j}]^{-1} 2^{jn_j} \beta_j^{n_j} = \sum_{\mathbf{v}} \gamma_s(\mathbf{v}) \prod_{j=1}^s \beta_j^{n_j}$$

since $\sum_{j=1}^s jn_j = s$.

□

3 The Moments of a Ratio of Two Random Variables

The next ingredient needed to prove our main result concerns the moments of a ratio of two random variables and is due, in essence, to SAWA [1972]. See also MEHTA and SWAMY ([1978], Lemma 3).

LEMMA 4: Let w_1 and w_2 be two random variables such that $P(w_2 > 0) = 1$. Assume that there exists a joint moment generating function of w_1 and w_2 :

$$\varphi(\theta_1, \theta_2) = E \exp(\theta_1 w_1 + \theta_2 w_2)$$

for all $|\theta_1| < \varepsilon$ and $-\infty < \theta_2 < \varepsilon$ where ε is some positive constant. Then, for $s = 1, 2, \dots$,

$$(8) \quad E(w_1/w_2)^s = \frac{1}{(s-1)!} \int_0^\infty t^{s-1} \left[\frac{\partial^s}{\partial \theta^s} \varphi(\theta, -t) \right]_{\theta=0} dt,$$

provided the expectation exists.

Proof: For $s = 1, 2, \dots$ we have

$$(9) \quad w_1^s = (\partial^s e^{\theta w_1} / \partial \theta^s)_{\theta=0}$$

and also

$$(10) \quad w_2^{-s} = \frac{1}{(s-1)!} \int_0^\infty t^{s-1} e^{-tw_2} dt.$$

$\left[\text{The latter equality, which holds for all } w_2 > 0, \text{ is obtained by substituting } \right.$
 $x = tw_2 \text{ in the gamma function } (s-1)! = \int_0^\infty x^{s-1} e^{-x} dx. \left. \right]$ Hence,

$$\begin{aligned}
 (11) \quad E(w_1/w_2)^s &= \frac{1}{(s-1)!} E \int_0^\infty t^{s-1} w_1^s \exp(-tw_2) dt \\
 &= \frac{1}{(s-1)!} \int_0^\infty t^{s-1} (E w_1^s \exp(-tw_2)) dt.
 \end{aligned}$$

where the first equality follows from (10) and the fact that $E(w_1/w_2)^s$ exists; the second equality is not completely trivial and follows from Fubini's theorem and an extension of it, known as Tonelli's theorem (see DUNFORD and SCHWARTZ [1958], section 3. 11). Also since all moments of w_1 exist it follows that $w_1^s \exp(\theta w_1)$ has a finite expectation and hence that $w_1^s \exp(\theta w_1 - tw_2)$ has a finite expectation. Thus

$$\begin{aligned}
 (12) \quad E w_1^s \exp(-tw_2) &= [E w_1^s \exp(\theta w_1 - tw_2)]_{\theta=0} \\
 &= \left[E \frac{\partial^s}{\partial \theta^s} \exp(\theta w_1 - tw_2) \right]_{\theta=0} \\
 &= \left[\frac{\partial^s}{\partial \theta^s} \varphi(\theta, -t) \right]_{\theta=0}.
 \end{aligned}$$

Inserting (12) in (11) gives the desired result. □

4 The Joint Moment Generating Function of s Quadratic Forms in Normal Variables

Before we can prove our main result we need one further lemma. This concerns the joint moment generating function of two quadratic forms in normal variables. Below we present the general case with s quadratic forms.

LEMMA 5 : Let x be a normally distributed $n \times 1$ vector with mean μ and positive definite covariance matrix $\Omega = LL'$. Let A_1, A_2, \dots, A_s be s symmetric $n \times n$ matrices. Then the joint moment generating function of $x' A_1 x, x' A_2 x, \dots, x' A_s x$:

$$\varphi(\theta_1, \theta_2, \dots, \theta_s) = E \exp(\theta_1 x' A_1 x + \dots + \theta_s x' A_s x)$$

exists for all $|\theta_i| < \varepsilon$ ($i = 1, 2, \dots, s$) where ε is some positive constant, and is given by

$$\varphi(\theta_1, \theta_2, \dots, \theta_s) = |I - 2C|^{-1/2} \exp\left(-\frac{1}{2} \eta' \eta\right) \exp\left(\frac{1}{2} \eta' (I - 2C)^{-1} \eta\right)$$

where

$$C = \theta_1 L' A_1 L + \dots + \theta_s L' A_s L, \quad \eta = L^{-1} \mu.$$

Proof: Let $u \cong N(0, I_n)$, so that $x = Lu + \mu$. Then

$$\begin{aligned} \varphi(\theta_1, \theta_2, \dots, \theta_s) &= E \exp(u + \eta)' C (u + \eta) \\ &= (2\pi)^{-(1/2)n} \int_{\mathbb{R}^n} \exp(u + \eta)' C (u + \eta) \cdot \exp\left(-\frac{1}{2} u' u\right) du \\ &= (2\pi)^{(1/2)n} \int_{\mathbb{R}^n} \exp -\frac{1}{2} \left\{ (u - \alpha)' (I - 2C) (u - \alpha) - 2R \right\} du \end{aligned}$$

where

$$\alpha = 2C(I - 2C)^{-1} \eta, \quad R = \frac{1}{2} \eta' (I - 2C)^{-1} \eta - \frac{1}{2} \eta' \eta.$$

Hence,

$$\begin{aligned} \varphi(\theta_1, \dots, \theta_s) &= e^R (2\pi)^{-(1/2)n} \int_{\mathbb{R}^n} \exp\left\{-\frac{1}{2} u' (I - 2C) u\right\} du \\ &= e^R |I_n - 2C|^{-1/2}, \end{aligned}$$

provided the θ 's are sufficiently close to zero to guarantee that $I_n - 2C$ is positive definite. \square

5 Main Result

We now have all the ingredients to prove our main result.

THEOREM 6 : Let x be a normally distributed $n \times 1$ vector with mean μ and positive definite covariance matrix $\Omega = LL'$. Let A be a symmetric

$n \times n$ matrix and B a positive semidefinite $n \times n$ matrix, $B \neq 0$. Let P be an orthogonal $n \times n$ matrix and Λ a diagonal $n \times n$ matrix such that

$$P' L' B L P = \Lambda, \quad P' P = I_n,$$

and define

$$A^* = P' L' A L P, \quad \mu^* = P' L^{-1} \mu.$$

Then we have, provided the expectation exists, for $s = 1, 2, \dots$,

$$\begin{aligned} E \left[\frac{x' A x}{x' B x} \right]^s &= \frac{\exp(-(1/2) \mu' \Omega^{-1} \mu)}{(s-1)!} \sum_v \gamma_s(v) \\ &\times \int_0^\infty t^{s-1} |\Delta| e^{(1/2) \xi' \xi} \prod_{j=1}^s (\text{tr } R^j + j \xi' R^j \xi)^{n_j} dt \end{aligned}$$

where the summation is over all $1 \times s$ vectors $v = (n_1, n_2, \dots, n_s)$ whose elements n_j are nonnegative integers satisfying $\sum_{j=1}^s j n_j = s$,

$$\gamma_s(v) = s! 2^s \prod_{j=1}^s [n_j! (2j)^{n_j}]^{-1}$$

and Δ is a diagonal positive definite $n \times n$ matrix, R a symmetric $n \times n$ matrix and ξ an $n \times 1$ vector given by

$$\Delta = (I_n + 2t\Lambda)^{-1/2}, \quad R = \Delta A^* \Delta, \quad \xi = \Delta \mu^*.$$

Proof: Let $\varphi(\theta, t)$ be the joint moment generating function of $x' A x$ and $x' B x$. Then, by Lemma 5

$$(13) \quad \varphi(\theta, -t) = |I - 2C|^{-1/2} \exp\left(-\frac{1}{2} \eta' \eta\right) \exp\left(\frac{1}{2} \eta' (I - 2C)^{-1} \eta\right)$$

where

$$C = \theta L' A L - t L' B L, \quad \eta = L^{-1} \mu.$$

Now, with P , Δ and R as defined in the theorem, we have

$$\begin{aligned} I - 2C &= I + 2t L' B L - 2\theta L' A L \\ &= P \Delta^{-1} (I - 2\theta R) \Delta^{-1} P', \end{aligned}$$

so that

$$(14) \quad |I - 2C|^{-1/2} = |\Delta|^{-1/2} |I - 2\theta R|^{-1/2}$$

and

$$(15) \quad (\mathbf{I} - 2\mathbf{C})^{-1} = \mathbf{P}\Delta(\mathbf{I} - 2\theta\mathbf{R})^{-1}\Delta\mathbf{P}'.$$

Letting $\xi = \Delta\mathbf{P}'\mathbf{L}^{-1}\mu$ and inserting (14), (15) and $\eta = \mathbf{L}^{-1}\mu$ in (13) gives

$$\begin{aligned} \varphi(\theta, -t) &= |\Delta| \cdot |\mathbf{I} - 2\theta\mathbf{R}|^{-1/2} \\ &\quad \times \exp\left(-\frac{1}{2}\mu'\Omega^{-1}\mu\right) \exp\left(\frac{1}{2}\xi'(\mathbf{I} - 2\theta\mathbf{R})^{-1}\xi\right) \\ &= \exp\left(-\frac{1}{2}\mu'\Omega^{-1}\mu\right) |\Delta| \exp\left(\frac{1}{2}\xi'\xi\right) \psi(\theta), \end{aligned}$$

where

$$\psi(\theta) \equiv |\mathbf{I} - 2\theta\mathbf{R}|^{-1/2} \exp\left(-\frac{1}{2}\xi'\xi\right) \exp\left(\frac{1}{2}\xi'(\mathbf{I} - 2\theta\mathbf{R})^{-1}\xi\right).$$

Now, let w be a normally distributed $n \times 1$ vector with mean ξ and covariance matrix \mathbf{I}_n . As a special case of Lemma 5 it appears that

$$\psi(\theta) = \mathbf{E} \exp(\theta w' \mathbf{R} w),$$

that is, $\psi(\theta)$ is the moment generating function of $w' \mathbf{R} w$. Thus, using the key property of moment generating functions,

$$\left[\frac{\partial^s \psi(\theta)}{\partial \theta^s} \right]_{\theta=0} = \mathbf{E} (w' \mathbf{R} w)^s.$$

Hence, since $\varphi(\theta, -t)$ depends on θ only through $\psi(\theta)$, we obtain

$$(16) \quad \left[\frac{\partial^s \varphi(\theta, -t)}{\partial \theta^s} \right]_{\theta=0} = \exp\left(-\frac{1}{2}\mu'\Omega^{-1}\mu\right) |\Delta| \exp\left(\frac{1}{2}\xi'\xi\right) \mathbf{E} (w' \mathbf{R} w)^s.$$

Also, from Lemma 3,

$$(17) \quad \mathbf{E} (w' \mathbf{R} w)^s = \sum_{\nu} \gamma_s(\nu) \prod_{j=1}^s \{\text{tr } \mathbf{R}^j + j \xi' \mathbf{R}^j \xi\}^{\nu_j}.$$

Inserting (17) in (16) and using Lemma 4, the result follows. □

6 Existence of the Moments of $x' \mathbf{A} x / x' \mathbf{B} x$

Theorem 1 is valid if and only if the expectation of $(x' \mathbf{A} x / x' \mathbf{B} x)^s$ exists. In this section we establish when this is the case.

THEOREM 7: Let x be a normally distributed $n \times 1$ vector with mean μ and positive definite covariance matrix $\Omega = LL'$. Let A be a symmetric $n \times n$ matrix and let B be a positive semidefinite $n \times n$ matrix of rank $r \geq 1$. If $r \leq n-1$, let Q be an $n \times (n-r)$ matrix of full column-rank $n-r$ such that $L'BLQ=0$.

(i) If $r \leq n-1$ and $L'ALQ=0$, or if $r=n$, then $E(x'A x/x'B x)^s$ exists for all $s \geq 0$.

(ii) If $r \leq n-1$, $Q'L'ALQ=0$ and $L'ALQ \neq 0$, then $E(x'A x/x'B x)^s$ exists for $0 \leq s < r$ and does not exist for $s \geq r$; and

(iii) If $r \leq n-1$ and $Q'L'ALQ \neq 0$, then $E(x'A x/x'B x)^s$ exists for $0 \leq s < r/2$ and does not exist for $s \geq r/2$.

Proof: Assume first that $r=n$, so that B is positive definite. Then

$$\frac{x'A x}{x'B x} = \frac{(B^{1/2} x)' B^{-1/2} A B^{-1/2} (B^{1/2} x)}{(B^{1/2} x)' (B^{1/2} x)},$$

which is bounded by the extreme eigenvalues of $B^{-1/2} A B^{-1/2}$, and hence possesses moments of any order.

Next assume $1 \leq r \leq n-1$. Let P_1 be an $n \times r$ matrix satisfying

$$L'BLP_1 = P_1 \Lambda_1, \quad P_1 P_1' = I_r, \quad P_1 Q = 0,$$

where Λ_1 is a diagonal $r \times r$ matrix containing the r positive eigenvalues of $L'BL$. Also let $P_2 = Q(Q'Q)^{-1/2}$ and define

$$y_1 = P_1 L^{-1} x \quad \text{and} \quad y_2 = P_2 L^{-1} x.$$

We note that y_1 and y_2 are independently normally distributed, and that

$$P_1 P_1' + P_2 P_2' = I_n.$$

Letting

$$C_{11} = P_1 L'ALP_1, \quad C_{12} = P_1 L'ALP_2, \quad C_{22} = P_2 L'ALP_2.$$

we obtain

$$(18) \quad \frac{x'A x}{x'B x} = \frac{y_1' C_{11} y_1 + 2y_1' C_{12} y_2 + y_2' C_{22} y_2}{y_1' \Lambda_1 y_1}.$$

It is clear from (18) that $E(x'A x/x'B x)^s$ exists if and only if

$$(19) \quad E \left[\frac{2y_1' C_{12} y_2 + y_2' C_{22} y_2}{y_1' \Lambda_1 y_1} \right]^s$$

exists. Now distinguish between three cases as in the theorem.

(i) $C_{12}=0, C_{22}=0$. Then (19) obviously exists for every s .

(ii) $C_{22}=0, C_{12} \neq 0$. Then (19) exists if and only if $E(y_1' C_{12} y_2 / y_1' \Lambda_1 y_1)^s$ exists. By Theorem 1 of KINAL [1980] this is the case iff $s < r$.

(iii) $C_{22} \neq 0$. Then (19) exists iff $E(y_1' y_1)^{-s}$ exists. But, since $y_1' y_1$ follows a noncentral χ^2 distribution with r degrees of freedom, $E(y_1' y_1)^{-s}$ exists iff $s < r/2$ (see JOHNSON and KOTZ [1969-1970], Chapter 26, section 2 and Chapter 30, section 3).

This completes the proof. □

7 Some Remarks on Computation

Let us define

$$(20) \quad f(v, t) = t^{s-1} |\Delta| e^{(1/2) \xi' \xi} \prod_{j=1}^s (\text{tr } R^j + j \xi' R^j \xi)^{n_j}$$

where Δ , R and ξ are defined in the theorem of section 5 and depend on t . The s -th moment of $x' A x / x' B x$ is then proportional to

$$(21) \quad \sum_v \gamma_s(v) \int_0^\infty f(v, t) dt.$$

The integral in (21) has to be evaluated numerically. If necessary, the entire integration range $(0, \infty)$ is first transformed to $(0, 1)$ using the identity

$$(22) \quad \int_0^\infty f(v, t) dt = \int_0^1 f\left(v, \frac{1-z}{z}\right) \frac{1}{z^2} dz.$$

Instead of (21) where a number of integrals have to be evaluated numerically, we can also calculate

$$(23) \quad \int_0^\infty \sum_v \gamma_s(v) f(v, t) dt.$$

The function under the integral sign is now more involved, but only one integral has to be evaluated. We found that (21) gives more accurate results than (23).

In addition, the expression

$$\text{tr } R^j + j \xi' R^j \xi$$

can be calculated either directly or indirectly by transforming R into diagonal form:

$$S'RS = M, \quad S'S = I_n$$

where S is orthogonal and M is diagonal. If μ_1, \dots, μ_n denote the diagonal elements of M and $\xi^* = S'\xi$, then

$$(24) \quad \text{tr } R^j + j \xi' R^j \xi = \sum_{i=1}^n \mu_i^j (1 + j \xi_i^{*2}).$$

We found that, unless n is small (say $n \leq 20$) and s is large (say $s \geq 8$), the direct approach is to be preferred. It is less time consuming and also more accurate.

A major simplification in the evaluation of the first four moments in the central case was proposed by ALI [1984], section 3, who noted that the product of factors involving t can be expressed as partial fractions.

Finally, as mentioned in section 2, there is, for a given value of s , the problem: how many vectors (n_1, n_2, \dots, n_s) are there? Let us designate this number as $p(s)$. Knowledge of $p(s)$ is important, if only to know how much memory space to reserve. It is easy to see that $p(s)$ is just the number of partitions of the integer s .² Values of $p(s)$ are given in HALL [1967], p. 35, for $s \leq 100$. For $s = 1, \dots, 12$ we find

s	1	2	3	4	5	6	7	8	9	10	11	12
$p(s)$	1	2	3	5	7	11	15	22	30	42	56	77

HALL [1967], p. 35, also provides a recurrence formula for $p(s)$ from which the values of $p(s)$ can be calculated for any value of s .

2. A partition of a positive integer s is a representation of s as a sum of positive integers $s = t_1 + t_2 + \dots + t_k$, $t_i > 0$, $i = 1, \dots, k$. For example, the partitions of the number 4 are (4), (3, 1), (2, 2), (2, 1, 1) and (1, 1, 1, 1), so that $p(4) = 5$.

●References

- ALI, M. M. (1984). — “Distributions of the Sample Autocorrelations when Observations are from a Stationary Autoregressive-Moving-Average Process”, *Journal of Business and Economic Statistics*, 2, pp. 271-278.
- CHAUBEY, Y. P. and NUR ENAYET TALUKDER, A. B. M. (1983). — “Exact Moments of a Ratio of Two Positive Quadratic Forms in Normal Variables”, *Communications in Statistics. Theory and Methods*, 12, pp. 675-679.
- DAVIS, A. W. (1979). — “Invariant Polynomials with Two Matrix Arguments Extending the Zonal Polynomials: Applications to Multivariate Distribution Theory”, *Annals of the Institute of Statistical Mathematics*, 31, part A, pp. 465-485.
- DEGOOYER, J. G. (1980). — “Exact Moments of the Sample Autocorrelations from Series Generated by General ARIMA Processes of Order (p, d, q) , $d=0$ or 1”, *Journal of Econometrics*, 14, pp. 365-379.
- DON, F. J. H. (1979). — “The Expectation of Products of Quadratic Forms in Normal Variables”, *Statistica Neerlandica*, 33, pp. 73-79.
- DUFOUR, J.-M. and ROY, R. (1985). — “Some Robust Exact Results on Sample Autocorrelations and Tests of Randomness”, *Journal of Econometrics*, 29, pp. 257-273.
- DUNFORD, N. and SCHWARTZ, J. T. (1958). — *Linear Operators (Part I: General Theory)*, Interscience Publishers, New York and London.
- DWIVEDI, T. D. and CHAUBEY, Y. P. (1981). — “Moments of a Ratio of Two Positive Quadratic Forms in Normal Variables” *Communications in Statistics. Simulation and Computation*, B10, pp. 503-516.
- GIDEON, R. A. and GURLAND, J. (1976). — “Series Expansions for Quadratic Forms in Normal Variables”. *Journal of the American Statistical Association*, 71, pp. 227-232.
- GURLAND, J. (1948). — “Inversion Formulae for the Distribution of Ratios”, *Annals of Mathematical Statistics*, 19, pp. 228-237.
- GURLAND, J. (1953). — “Distribution of Quadratic Forms and Ratios of Quadratic Forms”, *Annals of Mathematical Statistics*, 24, pp. 416-427.
- GURLAND, J. (1955). — “Distribution of Definite and Indefinite Quadratic Forms”, *Annals of Mathematical Statistics*, 26, pp. 122-127.
- GURLAND, J. (1956-1957). — “Quadratic Forms in Normally Distributed Random Variables”, *Sankhya*, 17, pp. 37-50.
- HALL, M. (1967). — *Combinatorial Theory*, Blaisdell Publishing Company, Waltham, Mass.
- HILLIER, G. and SMITH, M. (1983). — “On the Size of the Durbin-Watson Test Under Moderate Non-Normality”, *Discussion paper*, Département of Economics, Monash University.
- HOQUE, A. (1985). — “The Exact Moments of Forecast Error in the General Dynamic Model”, *Sankhya*, Series B, 47, pp. 128-143.
- HOQUE, A., and PETERS, T. A. (1986). — “Finite Sample Analysis of the ARMAX Models”, *Sankhya*, Series B, 48, to appear.
- IMHOF, P. J. (1961). — “Computing the Distribution of Quadratic Forms in Normal Variables”, *Biometrika*, 48, pp. 419-426.
- JAMES, A. T. (1964). — “Distributions of Matrix Variates and Latent Roots Derived from Normal Samples”, *Annals of Mathematical Statistics*, 35, pp. 475-501.

- JOHNSON, N. L. and KOTZ, S. (1969). — *Distributions in Statistics*, 3 vol., Houghton Mifflin Company, Boston.
- KENDALL, M. G. and STUART, A. (1977). — *The Advanced Theory of Statistics*. Vol. I, fourth edition, Charles Griffin & Co., London.
- KHURI, A. I. and GOOD, I. J. (1977). — “The Distribution of Quadratic Forms in Non-Normal Variables and an Application to the Variance Ratio”, *Journal of the Royal Statistical Society, Series B*, 39, pp. 217-221.
- KINAL, T. W. (1980). — “The Existence of Moments of k -Class Estimators”, *Econometrica*, 48, pp. 241-249.
- KUMAR, A. (1973). — “Expectation of Product of Quadratic Forms”, *Sankhya, Series B*, 35, pp. 359-362.
- LANCASTER, H. O. (1954). — “Traces and Cumulants of Quadratic Forms in Normal Variables”, *Journal of the Royal Statistical Society, Series B*, 16, pp. 247-254.
- LUGANNINI, R. and RICE, S. O. (1984). — “Distribution of the Ratio of Quadratic Forms in Normal Variables—Numerical Methods”, *SIAM Journal on Scientific and Statistical Computing*, 5, pp. 476-488.
- MAGNUS, J. R. (1978). — “The Moments of Products of Quadratic Forms in Normal Variables”, *Statistica Neerlandica*, 32, pp. 201-210.
- MAGNUS, J. R. (1979). — “The Expectation of Products of Quadratic Forms in Normal Variables: the Practice”, *Statistica Neerlandica*, pp. 131-136.
- MEHTA, J. S. and SWAMY, P. A. V. B. (1978). — “The Existence of Moments of Some Simple Bayes Estimators of Coefficients in a Simultaneous Equation Model”, *Journal of Econometrics*, 7, pp. 1-13.
- MORAN, P. A. P. (1948). — “Some Theorems on Time Series, II: The Significance of the Serial Correlation Coefficient”, *Biometrika*, 35, pp. 255-260.
- PHILLIPS, P. C. B. (1979). — “The Sampling Distribution of Forecasts from a First-Order Autoregression”, *Journal of Econometrics*, 9, pp. 241-261.
- SAWA, T. (1972). — “Finite-Sample Properties of the k -Class Estimators”, *Econometrica*, 40, pp. 653-680.
- SAWA, T. (1978). — “The Exact Moments of the Least Squares Estimator for the Autoregressive Model”, *Journal of Econometrics*, 8, pp. 159-172.
- SRIVASTAVA, V. K. and TIWARI, R. (1976). — “Evaluation of Expectations of Products of Stochastic Matrices”, *Scandinavian Journal of Statistics*, 3, pp. 135-138.
- TANEJA, V. S. (1976). — “Approximations to the Distribution of Indefinite Quadratic Forms in Noncentral Normal Variables”, *Metron*, 34, pp. 255-268.
- WALKER, J. J. (1979). — “An Asymptotic Expansion for Unbalanced Quadratic Forms in Normal Variables”, *Journal of the American Statistical Association*, 74, pp. 389-392.