

Testing for Super-Exogeneity in the Presence of Common Deterministic Shifts

Hans-Martin KROLZIG, Juan TORO *

ABSTRACT. – This paper introduces the concept of common deterministic shifts (CDS). This concept is simple, intuitive and relates to the common structure of shifts or policy interventions. We propose a Reduced Rank technique to investigate the presence of CDS. The proposed testing procedure has standard asymptotics and good small-sample properties. We further link the concept of CDS to that of super-exogeneity. It is shown that CDS tests can be constructed which allow to test for super-exogeneity. The Monte Carlo evidence indicates that the CDS test for super-exogeneity dominates testing procedures proposed in the literature.

Test de super-exogénéité en présence de rupture déterministe commune

RÉSUMÉ. – Cet article introduit le concept de ruptures déterministes communes (RDC). Ce concept simple et intuitif est relié à des déformations de structures communes ou à des interventions politiques. Nous proposons d'utiliser une technique de rang réduit pour étudier la présence de RDC. La procédure de test considérée a un comportement asymptotique standard et de bonnes propriétés à distance finie. Nous relierons de plus le concept de RDC au problème de la superexogénéité. Nous montrons que nous pouvons utiliser des tests de RDC pour tester la super-exogénéité. Un exercice de Monte-Carlo semble indiquer que les tests RDC pour la super-exogénéité dominent les procédures de test existantes.

* H.M. KROLZIG: Department of Economics and Nuffield College, Oxford;
J. TORO: European University Institute, Florence and Centra (Seville).

We are grateful to the guest editor, S. GREGOIR, and two anonymous referees for valuable comments and suggestions on a previous version of this paper. We also thank D. HENDRY, S. JOHANSEN, U. HASSLER and B. NIELSEN for useful comments and discussions. Financial support from the UK Economic and Social Research Council under grant L138251009 is gratefully acknowledged by the first author.

1 Introduction

Deterministic shifts in the conditional mean of economic variables are a recurrent feature in empirical economics. These shifts happen to affect not just one single economic variable but, contemporaneously, also other related economic variables. Furthermore, some economic time series are affected by several shifts. These shifts might be related linearly and this linear relationship might prevail throughout time. We will refer to these shifts as common deterministic shifts. Here we propose a technique that can be used to analyze such phenomena, and can help to gather important information about how breaks are related through economic variables and across time.

Frequently, deterministic shifts are induced by policy changes. Policymakers move the level of some variables in order to affect some target variables and reach specific goals. When deterministic shifts are induced by policymakers, the relationship between common deterministic shifts and super-exogeneity becomes apparent. Super-exogeneity (see ENGLE, HENDRY and RICHARD [1983]) establishes conditions under which the parameters of the partial model are invariant to changes in the parameters of the marginal model. In an economic context, the marginal model can be thought of as the instrument of the policymaker (*say*, the interest rate). The partial model could be thought of as the process for the goal variable (*say*, the rate of inflation). Super-exogeneity sets conditions under which the partial model has invariant parameters, which allows its use for policy analysis despite changes in the marginal model. The concepts of common deterministic shifts and super-exogeneity are closely related if we limit the set of policymakers interventions to changes in the conditional mean of the marginal process (*say*, the level of interest rates or the rate of growth of money).

In this article, we introduce the concept of common deterministic shifts (CDS) and propose how they can be estimated and tested. The proposed testing procedure possesses standard asymptotics and the small sample properties are investigated with a Monte Carlo study. The Monte Carlo study shows that the test for common deterministic shifts has good size and power properties in small samples.

We also analyze the relationship between common deterministic shifts and super-exogeneity. We show how a test for super-exogeneity could be implemented as a set of restrictions in the reduced rank matrices of the intervention variables used to model the CDS. The empirical applicability of the alternative super-exogeneity test that we propose is exemplified with the Lucas supply function (LUCAS [1975]). Three different testing procedures are introduced to test for super-exogeneity within the framework of common deterministic shifts. The first test reparameterizes the original approach of ENGLE and HENDRY [1993] as a reduced rank restriction. In this case the null corresponds to that proposed in ENGLE and HENDRY [1993], but the alternative is defined differently. The other two tests proposed differ in how the model is estimated under the null. A Monte Carlo experiment concludes that for a given size, the CDS test for super-exogeneity has a higher power than the ENGLE and HENDRY [1993] test. Moreover, we further discuss how the empi-

rical applicability of the idea of CDS does not have to be restricted to the analysis of super-exogeneity and we present other examples where this methodology could be useful. We also discuss how this idea could be implemented in the case where $I(1)$ variables are modelled and also in the situation in which the dates of the shifts are unknown.

The structure of the paper is as follows: In the next section, we introduce the concept of common deterministic shifts. In Section 3, we define the model and introduce a reduced rank technique to estimate and test for common deterministic shifts. The size and power of the proposed technique are investigated in Section 4 with a Monte Carlo simulation experiment. Section 5 discusses the concept of super-exogeneity in the framework of common deterministic shifts. In Section 6, we develop testing procedures for super-exogeneity, which are based on the existence of common deterministic shifts. Their small-sample properties are analyzed in a small Monte Carlo experiment discussed in Section 7. The empirical applicability of the introduced tests is illustrated in Section 8, which also comments on their limitations and possible extensions. Last, Section 9 concludes.

2 The Concept of Common Deterministic Shifts

ENGLE and KOZICKI [1993] have recently proposed the idea of common features in time series. This idea is inspired by the concept of cointegration introduced in ENGLE and GRANGER [1987]. ENGLE and KOZICKI [1993] show that a feature is common to a set of time series if a linear combination of them does not have the feature though each of the series individually has it. Some particular examples of this concept are the idea of common cycles, introduced by ENGLE and KOZICKI [1993], and the idea of co-breaking, introduced by HENDRY [1997]. The concept of co-breaking is closely related to the idea of cointegration: while cointegration removes unit roots from linear combinations of variables, co-breaking can eliminate the effects of regime shifts by taking linear combinations of variables.

DEFINITION 1: Consider $\{x_t\}$ to be an n dimensional vector process, which is modeled as a vector autoregressive (VAR) process of order p , $A(L)x_t = \mu_t + \varepsilon_t$. Consider a set of $n \leq s < T$ dummy variables $d_{t,i}$ each of which is zero except for unity at times $t \in \mathcal{T}_i$, such that $\mu_t = \sum_{i=1}^s \mu_i d_{t,i}$, or $\mu_t = \mathcal{M}D_t$, where \mathcal{M} is $n \times s$ and D_t is $s \times 1$ with $D_t = (d_{t,1} : d_{t,2} : \dots : d_{t,s})'$. The condition for common deterministic shifts can be written as $\Phi' \mathcal{M} = \mathbf{0}$. Thus we have that $r = \text{rank} [\mathcal{M}] < n$ is necessary and sufficient for $\Phi' \mu_t = 0$ for all $t \in \mathcal{T}$, where $\Phi \neq \mathbf{0}$ is $n \times (n - r)$. We say that the equations in the VAR are subject to common deterministic shifts (CDS) if shifts taking place across the n individual equations are linearly related.

In the definition of common deterministic shifts we just require that shifts be related across variables and through time, which can be expressed as a convenient reduced rank condition in the coefficients of the intervention variables. This concept is different from the co-breaking concept of HENDRY [1997] which requires that linear combinations of variables cancel the shifts in the processes. In the following, we consider the n -dimensional linear Gaussian VAR(p):

$$(1) \quad x_t = \sum_{i=1}^p A_i x_{t-i} + \mu_t + \varepsilon_t,$$

where $\varepsilon_t \sim \text{NID}(\mathbf{0}, \Sigma)$ and the roots of the vector autoregressive polynomial are within the unit circle, $\left| \mathbf{I} - \sum_{i=1}^p A_i z^i \right| = 0 \implies |z| > 1$. Thus there are no unit roots in the system and the possible non-stationarity is due to the deterministic breaks. This implies that the process possesses the infinite-order vector moving-average representation:

$$x_t = \sum_{i=0}^{\infty} \Psi_i (\varepsilon_{t-i} + \mu_{t-i}),$$

where $A(L)\Psi(L) = \mathbf{I}$, and in the case of a VAR(1) we have that $\Psi_i = A^i$.

The methodology proposed in this paper relies on using appropriate shift dummies for known dates. It is assumed that the points at which these shifts occur are known, which avoids the problem of some nuisance parameters. In order to illustrate the reduced rank approach consider a VAR(1) with shifts in the intercept:

$$(2) \quad x_t = A_1 x_{t-1} + \sum_{i=1}^s \mu_i d_{t,i} + \varepsilon_t.$$

Furthermore, suppose that the shifts are permanent. Then we can use the corresponding shift dummies to model them: $d_{t,i} = I(t > t_i)$, where $I(\bullet)$ is the indicator function and $1 < t_i < T$.

CDS is at least of order $(n - r)$ if there exist $(n - r)$ linearly independent vectors ϕ_i satisfying:

$$\phi_i' \mu_t = \phi_i' \mathcal{M} D_t = 0$$

such that the $n \times (n - r)$ matrix $\Phi = (\phi_1 : \dots : \phi_{(n-r)})$ has rank $(n - r)$. Then, $\Phi' \mathcal{M} = \mathbf{0}$ so rank $(\mathcal{M}) < n$ and the nullity of \mathcal{M} determines the order of CDS. In other words, CDS implies that \mathcal{M} is of reduced rank. Thus, \mathcal{M} can be decomposed into the product of two matrices, η and ξ of full rank r , such that we can rewrite (2) as:

$$(3) \quad x_t = A_1 x_{t-1} + \eta \xi' D_t + \varepsilon_t.$$

Note that the matrices η and ξ are not unique without suitable normalization: if H is any $r \times r$ non-singular matrix, then $\mathcal{M} = \eta\xi'$ implies that $\eta HH^{-1}\xi' = \eta H(\xi H^{-1})' = \eta^*\xi^{*'} = \mathcal{M}$ as well. If common deterministic shifts are a property of the data, the coefficient matrix of the dummy regressors will have a reduced rank.

3 Estimating CDS Vectors by Reduced Rank Regressions

3.1 The Reduced-Rank Regression Problem

Maximum likelihood estimation of the $\text{CDS}(n-r) - \text{VAR}_n(p)$ is close to the analysis of the likelihood in cointegrating systems, and both are based on the reduced rank regression technique introduced by ANDERSON [1958] and Tso [1981]. The analogy with the cointegration model is straightforward if one bears in mind that the regime-dummies d_t behave like non-stationary processes if there are structural breaks. In this case the matrix \mathcal{M} determines how the non-stationarity feeds into the variables of the system: the rank r of the matrix \mathcal{M} gives the number of *common deterministic shifts*, and the CDS rank ($n-r$ or $s-r$) gives the dimension of the space whose one-step predictions are free from these *deterministic breaks*. In contrast to the cointegration problem, the number of breaks s is not necessarily identical to the number of endogenous variables in the system, such that the matrix \mathcal{M} is $n \times s$ with rank $r \leq \min(n, s)$.

In matrix notation, we have:

$$(4) \quad X = \mathcal{B}Z + \mathcal{M}D + E,$$

where $X := (x_1 : x_2 : \dots : x_T)$ is $n \times T$, $Z := (z_1 : z_2 : \dots : z_T)$ is $pn \times T$ with $z'_t := (x'_{t-1} : \dots : x'_{t-p})$, $\mathcal{B} := (A_1 : \dots : A_p)$ is $n \times pn$, $E = (\varepsilon_1 : \varepsilon_2 : \dots : \varepsilon_T)$ is $n \times T$ and D is $s \times T$.¹

\mathcal{M} is $n \times s$ and of rank r and can be decomposed in two full rank matrices, $\mathcal{M} = \eta\xi'$, where η are the loadings and ξ the linear relationships across breaks.

The log-likelihood function for a sample size T is easily seen to be:

$$(5) \quad \ln L = -\frac{nT}{2} \ln 2\pi - \frac{T}{2} \ln |\Sigma| - \frac{1}{2} \text{tr} \left[(X - \mathcal{B}Z - \eta\xi'D)' \Sigma^{-1} (X - \mathcal{B}Z - \eta\xi'D) \right].$$

1. Note that, for reasons of simplicity, we assume here that the variables have a zero mean before the first break. If the mean of the variables is non-zero before the first break, an intercept ν can be added to the model, such that $z'_t := (1 : x'_{t-1} : \dots : x'_{t-p})$ and $\mathcal{B} := (\nu : A_1 : \dots : A_p)$ is $n \times (1 + pn)$.

3.2 Estimation of \mathcal{B} and Σ Conditional on \mathcal{M}

Note that for any fixed η and ξ , the maximum of $\ln L$ is obtained by:

$$(6) \quad \tilde{\mathcal{B}}(\eta\xi') = (X - \eta\xi'D)Z'(ZZ')^{-1}.$$

If we substitute \mathcal{B} in (5) by (6), we get:

$$(7) \quad \ln L = -\frac{nT}{2}\ln 2\pi - \frac{T}{2}\ln |\Sigma| - \frac{1}{2}\text{tr} \left[(XH - \eta\xi'DH)' \Sigma^{-1} (XH - \eta\xi'DH) \right],$$

where $H := (\mathbf{I}_T - Z'(ZZ')^{-1}Z)$ is a $T \times T$ matrix. Hence, we just have to maximize this expression with respect to η and Σ . For given η and ξ , the maximum is obtained for:

$$\tilde{\Sigma}(\eta\xi') = T^{-1}(X - \eta\xi'D)H(X - \eta\xi'D)'$$

Consequently we must maximize:

$$(8) \quad -\frac{T}{2}\ln \left| T^{-1}(X - \eta\xi'D)H(X - \eta\xi'D)' \right|$$

or, equivalently, minimize the determinant with respect to η and ξ (see LÜKEPOHL [1991]).

3.3 Estimation of η Conditional on ξ

Note that in (7), X and D are corrected for Z . Define the corresponding residuals as:

$$\begin{aligned} R_X & : = XH, \\ (n \times T) & \\ R_D & : = DH, \\ (s \times T) & \end{aligned}$$

and the corresponding moment matrices as:

$$S_{ij} = T^{-1}R_i R_j' \text{ for } i, j = X, D.$$

Then (8) can be rewritten as:

$$(9) \quad -\frac{T}{2}\ln \left| T^{-1}(R_X - \eta\xi'R_D)(R_X - \eta\xi'R_D)' \right|.$$

For fixed ξ , (8) is maximized with respect to matrix η by regression:

$$(10) \quad \begin{aligned} \tilde{\eta}(\xi) &= R_X (\xi'R_D)' [(\xi'R_D)(\xi'R_D)']^{-1} \\ &= R_X R_D' \xi [\xi'(R_D R_D')\xi]^{-1} \\ &= S_{XD} \xi (\xi' S_{DD} \xi)^{-1}. \end{aligned}$$

3.4 Estimation of ξ

Apart from a constant, the concentrated log-likelihood for our reduced rank problem can be shown to be:

$$\begin{aligned}
 (11) \quad & \ln L \left(\tilde{\eta}(\xi), \xi, \tilde{\Sigma}(\tilde{\eta}(\xi), \xi') \right) \\
 &= -\frac{T}{2} \ln \left| \tilde{\Sigma}(\tilde{\eta}(\xi), \xi') \right| \\
 &= -\frac{T}{2} \ln \left| \frac{1}{T} (R_X - R_X R_D' \xi [\xi' (R_D R_D') \xi]^{-1} \xi' R_D) (R_X - R_X R_D' \xi [\xi' (R_D R_D') \xi]^{-1} \xi' R_D)' \right| \\
 &= -\frac{T}{2} \ln \left| \frac{1}{T} R_X (\mathbf{I} - R_D' \xi [\xi' (R_D R_D') \xi]^{-1} \xi' R_D) (\mathbf{I} - R_D' \xi [\xi' (R_D R_D') \xi]^{-1} \xi' R_D)' R_X' \right| \\
 &= -\frac{T}{2} \ln \left| \frac{1}{T} R_X \left[\mathbf{I} - (\xi' R_D)' \left[(\xi' R_D) (\xi' R_D) \right]^{-1} (\xi' R_D) \right] R_X' \right| \\
 &= -\frac{T}{2} \ln |S_{XX} - S_{XD} \xi (\xi' S_{DD} \xi)^{-1} \xi' S_{DX}|.
 \end{aligned}$$

By using the identity:

$$\begin{aligned}
 \left| \begin{array}{cc} S_{XX} & S_{XD} \xi \\ \xi' S_{DX} & \xi' S_{DD} \xi \end{array} \right| &= |\xi' S_{DD} \xi| \left| S_{XX} - S_{XD} \xi (\xi' S_{DD} \xi)^{-1} \xi' S_{DX} \right| \\
 &= |S_{XX}| \left| \xi' S_{DD} \xi - \xi' S_{DX} S_{XX}^{-1} S_{XD} \xi \right|,
 \end{aligned}$$

Equation (11) can be expressed as:

$$\begin{aligned}
 \ln L \left(\tilde{\eta}(\xi), \xi, \tilde{\Sigma}(\tilde{\eta}(\xi), \xi') \right) &= -\frac{T}{2} \ln \frac{|S_{XX}| \left| \xi' S_{DD} \xi - \xi' S_{DX} S_{XX}^{-1} S_{XD} \xi \right|}{|\xi' S_{DD} \xi|} \\
 &= -\frac{T}{2} \ln \frac{|S_{XX}| \left| \xi' (S_{DD} - S_{DX} S_{XX}^{-1} S_{XD}) \xi \right|}{|\xi' S_{DD} \xi|}.
 \end{aligned}$$

Hence, the maximum of $\ln L$ is given by:

$$\min_{\xi} \frac{\left| \xi' (S_{DD} - S_{DX} S_{XX}^{-1} S_{XD}) \xi \right|}{|\xi' S_{DD} \xi|}$$

and following a basic theorem of matrix analysis (see, for example, JOHANSEN [1995], Lemma A.8), this factor is minimized among all $n \times r$ matrices ξ by solving the eigenvalue problem:

$$\left| \rho S_{DD} - (S_{DD} - S_{DX} S_{XX}^{-1} S_{XD}) \right| = 0$$

or, for $\lambda = 1 - \rho$, the eigenvalue problem:

$$\left| \lambda S_{DD} - S_{DX} S_{XX}^{-1} S_{XD} \right| = 0$$

for eigenvalues λ_i and eigenvectors v_i , such that:

$$(12) \quad \lambda_i S_{DD} v_i = S_{DX} S_{XX}^{-1} S_{XD} v_i.$$

If we normalize ξ such that $\xi' S_{DD} \xi = \mathbf{I}_r$ then the vectors of $\tilde{\xi}$ are given by the eigenvectors corresponding to the r largest eigenvalues of $S_{DD} - S_{DX} S_{XX}^{-1} S_{XD}$. The maximum log-likelihood under the rank (\mathcal{M}) = r restriction is given by:

$$(13) \quad \max \ln L = -\frac{nT}{2} \ln 2\pi - \frac{T}{2} \left[\ln |S_{XX}| + \sum_{i=1}^r \ln (1 - \hat{\lambda}_i) \right] - \frac{nT}{2},$$

since, by choice of $\tilde{\xi}$, we have that $\xi' S_{DD} \xi = \mathbf{I}_r$, as well as $\xi' S_{DX} S_{XX}^{-1} S_{XD} \xi = \text{diag}(\hat{\lambda}_1, \dots, \hat{\lambda}_r)$, where $\hat{\lambda}_1, \dots, \hat{\lambda}_r$ are the r largest eigenvalues solutions of (12).

3.5 Testing for the CDS Rank

Since $\text{CDS}(n-r)$ implies $\text{CDS}(n-r-1)$, it seems natural to seek the maximum degree of CDS. In general, two cases have to be distinguished. In the first case, the number of potential breaks s is less than the dimension of the system n , $m = \min(s, n) = s < n$. In the second case, the number of potential breaks s is not less than the dimension of the system n , *ie*, $m = \min(s, n) = n \leq s$. Following ANDERSON [1958], the likelihood ratio test statistic for testing the $\text{CDS}(r)$ against the $\text{CDS}(n)$ is given by:

$$-2 \ln Q(H(r) | H(n)) = -T \sum_{i=r+1}^n \ln (1 - \hat{\lambda}_i),$$

which has a χ^2 -distribution with degrees of freedom equal to $(n-r)(s-r)$.

Suppose in the following that $n \leq s$. Then, the following hypotheses might be of interest:

- (i) $\text{CDS}(n)$: $\text{rank}(\mathcal{M}) = 0$. No breaks. We would calculate the statistic $Q_0 = -2 \log Q(H(0) | H(1))$ and compare it with the relevant quantile of the corresponding χ^2 -distribution.
- (ii) $\text{CDS}(n-r)$: $\text{rank}(\mathcal{M}) = r$, $0 < r < n$. There are breaks which are common to the processes. In this case we would calculate the statistic $Q_r = -2 \log(H(r) | H(r+1))$ and compare it with the relevant quantile.

If we had no prior knowledge about r one could use the same testing logic proposed by JOHANSEN [1995] to determine the cointegrating rank. First, we would compare Q_0 with its quantile, say, c_0 . If $Q_0 < c_0$, we take $\hat{r} = 0$. If $Q_0 \geq c_0$, we would calculate Q_1 and compare it with its quantile, c_1 . If $Q_1 < c_1$, we take $\hat{r} = 1$, otherwise, if $Q_1 \geq c_1$ we would continue as before calculating sequentially the corresponding test and comparing it with the appropriate quantile. This testing procedure defines an estimator \hat{r} with values $0, 1, \dots, n$ which converges in probability to the true value.

4 Some Monte Carlo Results

4.1 Design of the Monte Carlo

In this section we analyze the size and power of the rank test for common deterministic shifts. The data generation process (DGP) is a two-dimensional process $x_t = (y_t : z_t)'$ where z_t does not feedback into the first equation and there are two breaks in the intercept term at time t_1 and t_2 ,

$$(14) \quad \begin{pmatrix} y_t \\ z_t \end{pmatrix} = \begin{pmatrix} 0.75 & 0.5 \\ 0 & \alpha \end{pmatrix} \begin{pmatrix} y_{t-1} \\ z_{t-1} \end{pmatrix} + \mathcal{M} \begin{pmatrix} d_{t_1} \\ d_{t_2} \end{pmatrix} + \begin{pmatrix} \varepsilon_{1t} \\ \varepsilon_{2t} \end{pmatrix}$$

where $\begin{pmatrix} \varepsilon_{1t} \\ \varepsilon_{2t} \end{pmatrix} \sim \text{NID} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right)$.

The matrix of intervention coefficients \mathcal{M} embeds information about the size of the shifts, and the relationship of the shifts across variables and time. The interpretation of the model will critically depend on the assumed rank of the matrix \mathcal{M} :

- (i) $\text{rank}(\mathcal{M}) = 0$. No breaks.
- (ii) $\text{rank}(\mathcal{M}) = 1$. There are breaks which are common to both processes.
- (iii) $\text{rank}(\mathcal{M}) = 2$ (full rank). There are breaks independent to each process.

We consider two specifications for \mathcal{M} . In the first specification, \mathcal{M} is of full rank:

$$\mathcal{M}_1 = k \begin{pmatrix} 1 & .5 \\ .5 & 1 \end{pmatrix}.$$

The second specification for \mathcal{M} , denoted \mathcal{M}_2 , has reduced rank:

$$\mathcal{M}_2 = k \begin{pmatrix} 1 \\ \varphi \end{pmatrix} (1 \quad \pi).$$

This corresponds to the case of CDS, where φ defines the relationship of the breaks across equations, π defines the relationship of breaks across time and k defines the magnitude of the breaks measured in terms of the standard deviation. In the Monte Carlo experiment, we analyze the properties of the CDS test when we allow φ , π and k to vary. We will consider the following values: $\varphi \in \{0.25, 0.5, 1\}$, $\pi \in \{0.25, 0.5, 1\}$ and $k \in \{1, 1.5, 2\}$.

The benchmark case will have $\alpha = 0.8$, but one might be interested in analyzing the size of the test when the process for z_t becomes close to the unit root. That is, $\alpha \in \{0.95, 0.975, 0.99\}$. Furthermore, in order to study how the distance between breaks affects the CDS test, we consider different distances

between the breaks. As the break points are defined as fractions of the sample size, $t_1 = \tau_1 T$ and $t_2 = \tau_2 T$, we will parametrically vary the first break point, $\tau_1 \in \{0.30, 0.31, \dots, 0.6\}$, while keeping the second break point fixed, $\tau_2 = 0.7$.

We consider three different sample sizes: $T = 50, 100$ and 150 . The number of replications is $N = 10000$. If we had relied on a full factorial design, the number of experiments would have been 18225. We therefore focus on two benchmark DGPs in our Monte Carlo study: DGP_1 is characterized by $(\mathcal{M}_1; k = 2; \alpha = 0.8; \tau_1 = 0.3, \tau_2 = 0.7)$ and will be used to study the power properties of the CDS test. DGP_2 is defined by $(\mathcal{M}_2; \varphi = 1, \pi = 1, k = 2; \alpha = 0.8; \tau_1 = 0.3, \tau_2 = 0.7)$ and allows to investigate the size properties of the CDS test given the reduced rank of its matrix of intervention coefficients.

4.2 Evaluation of the Monte Carlo

Table 1 presents the rejection frequencies of the CDS test for our simulated DGP_2 for nominal sizes from 1% to 20%. It can be seen that the real size differs from the nominal size and this difference cannot be explained by Monte Carlo sampling variation. Table 2 presents the rejection frequencies for the test statistic $H_0 : \text{rank}(\mathcal{M}) = 1$ against $H_1 : \text{rank}(\mathcal{M}) = 2$ when the model simulated is DGP_1 . This delivers the power of the CDS test for our benchmark case.

Figure 1 compares the theoretical and empirical quantiles of the CDS test statistic for the reduced rank of \mathcal{M} : $H_0 : \text{rank}(\mathcal{M}) = 1$ against $H_1 : \text{rank}(\mathcal{M}) = 2$. The simulated model is the benchmark DGP_2 and the sample sizes are $T = 50, 100$ and 150 . The distribution of the test follows asymptotically a χ^2 with one degree of freedom. The results indicate that the empirical size is slightly higher than the nominal size, but the simulated quantiles approach the theoretical quantiles of the asymptotic distribution with increasing sample size relatively fast for $T \geq 50$.

In order to compare the power of the CDS test against its true size we will make use of size-power curves. We thus construct two different experiments: in the first experiment the DGP satisfies the null of common deterministic shifts, and in the second experiment the matrix of interventions coefficients has full rank.

TABLE 1
Size of the CDS Rank Test

T	20%	10%	5%	1%
50	0.263	0.156	0.088	0.0218
100	0.234	0.128	0.066	0.0146
150	0.230	0.118	0.061	0.0136

Note: Rejection frequencies for the test statistic $H_0 : \text{rank}(\mathcal{M}) = 1$ against $H_1 : \text{rank}(\mathcal{M}) = 2$ for the benchmark DGP_2 .

TABLE 2
Power of the CDS Rank Test

T	20%	10%	5%	1%
50	0.722	0.564	0.413	0.169
100	0.910	0.806	0.665	0.343
150	0.968	0.909	0.805	0.469

Note: Rejection frequencies for the test statistic $H_0 : \text{rank}(\mathcal{M}) = 1$ against $H_1 : \text{rank}(\mathcal{M}) = 2$ for the benchmark DGP_1 .

FIGURE 1
QQ Plot of the CDS Test for the Reduced Rank of \mathcal{M}





FIGURE 2
Size-Power Curve Analysis



Let us denote the test statistic in replication j by ϑ_j . With ϑ_j there is an associated p value ($p_j = p(\vartheta_j)$), obtained from our asymptotic distribution, which is $\chi^2(1)$. For any point x_i in the $(0, 1)$ interval we can define the empirical distribution function (EDF) of p_j as:

$$F(x_i) = \frac{1}{N} \sum_{j=1}^N I(p_j < x_i).$$

FIGURE 3
Size-Power Curve Analysis for the CDS Rank Test





FIGURE 4
Size of the Test Statistic for Different Values of α and τ_1 Varying from 0.3 to 0.6



The EDF $F(x_i)$ reports the probability of getting a p value less than x_i under the null. Following the suggestion of DAVIDSON and MACKINNON [1997] the EDF is evaluated at m points x_i , $i = 1, \dots, m$, where $m = 215$ and $x_i \in \{0, 0.001, 0.002, \dots, 0.999\}$. We can then obtain a p -value plot plotting $F(x_i)$ against x_i . If the test is well-behaved (*i.e.*, the test is rejected just about the right proportion of times) the plot should be close to the 45 degree line.

If we denote by $F^*(x_i)$ the EDF under the alternative, and the same random numbers are used when simulating the errors under the null and the alternative, then the size-power curve is obtained plotting $F^*(x_i)$ against $F(x_i)$.

The p -value plots in Figure 2 help to summarize and complement the information contained in Tables 1 and 2 and Figure 1. In the left panel of Figure 1, a p -value plot of the test statistic of $H_0: \text{rank}(\mathcal{M}) = 1$ against $H_1: \text{rank}(\mathcal{M}) = 2$ is plotted which is based on the empirical distribution of the p -values of our test statistic when the simulated DGP is the benchmark DGP_2 . We can see that the CDS test tends to overreject slightly. This overrejection tends to decrease quickly as the sample size increases with the p -value plot approaching the 45 degree line. The plot is truncated for values of $x_i > 0.2$ since relative small test sizes are of interest. The right panel of Figure 2 depicts the corresponding power-size curve. The power always exceeds hugely the actual size and is excellent for $T = 100$ and 150.

The robustness of the previous results is analyzed in Figure 3, which presents evidence of the size-power trade-off under parametric variations of the benchmark DGP_2 . Each plot contains the size-power curve (bold line) for the CDS test statistic for the presence of one common deterministic shift, $H_0: \text{rank}(\mathcal{M}) = 1$ against $H_1: \text{rank}(\mathcal{M}) = 2$ for three sample sizes $T = 50, 100$ and 150. For each parametrisation of the DGP, the size-power curve is compared with the size-power curve in the benchmark DGP_2 (thin line).


Plots [A] and [B] compare the size-power curves in the benchmark DGP_2 with $k = 1.5$ and $k = 1$ respectively. Plot [C] compares the size-power curves in the benchmark DGP_2 with that obtained when $\varphi = 0.5$. Plot [D] visualizes the effects of $\pi = 0.5$. Finally, Plot [E] graphs the size-power curve obtained when $k = 1$ and $\varphi = \pi = 0.5$. From the plots, we can see that the power for a given size remains largely unchanged with different parametrisation of the benchmark DGP. There is, however, a substantial reduction in power relative to our benchmark DGP_2 when the size of the breaks is lowered as in the case of $k = 1$ and $\varphi = \pi = 0.5$.

We next analyze how the position of the breaks may affect the size and the power of the CDS test by changing the point of occurrence of the first break, allowing τ_1 to vary from 0.3 to 0.6 of the sample ($\tau_1 \in \{0.30, 0.31, \dots, 0.6\}$). The second break is always fixed at $t_2 = \tau_2 T$ for $\tau_2 = 0.7$. The resulting size and power is analyzed for four different values of α in the benchmark DGPs , $\alpha \in \{0.8, 0.95, 0.975, 0.99\}$, which allows for the investigation of the behavior of the test as the process for z_t gets closer to a unit root process. As in the previous experiments, three different sample sizes ($T = 50, 100$ and 150) are considered.

The Monte Carlo results shown in Figures 4 and 5 depict the size and power of the CDS test for a given nominal size of 5%. In these plots the vertical

FIGURE 5

Power of the Test Statistic for Different Values of α and τ_1 Varying from 0.3 to 0.6



axis measures the size and the power, respectively, and the horizontal axis measures the location of the first break as a fraction of the sample size. When interpreting the plots, the reader should be aware that Figure 5 depicts the power for the given *nominal* size. From the graphs one can see that the actual size clearly deviates from the nominal size when α gets closer to unity. Furthermore, for $\alpha = .8$ the actual size increases rapidly as the break points get closer to the second break point. For $T = 50$ power increases with greater distance from the second break. The power is uniformly strong for $T \geq 100$. Overall, the Monte Carlo results are very promising. The asymptotic χ^2 distribution of the CDS test is found to provide a good approximation of the small-sample properties of the test.

5 Super-Exogeneity in the Presence of Common Deterministic Shifts

The previous section used reduced rank regressions as a modelling device for finding common deterministic shift features. In this section, we present an alternative analysis. Though the concept of common deterministic shifts has been introduced in terms of an unrestricted model, its most insightful application involves conditional models.

5.1 The Concept of Super-Exogeneity

Before we discuss how common deterministic shifts in the conditional mean are related to the concept of super-exogeneity, let us introduce the definitions of weak and super-exogeneity. We will consider a statistical model for the vector process $x_t = (y_t' : z_t')'$ with parameters $\theta \in \Theta$. We are interested in a function of θ , called the parameters of interest, *ie*, $\vartheta = \vartheta(\theta)$.

DEFINITION 2: *Weak exogeneity: The process z_t is called weakly exogenous for the parameter of interest ϑ if there exist a parametrisation of the model such that:*

$$f(x_1, \dots, x_T; \alpha, \beta) = \prod_{t=1}^T g_t(y_t | z_t, x_1, \dots, x_{t-1}; \beta) h_t(z_t | x_1, \dots, x_{t-1}; \alpha),$$

where $(\alpha, \beta) \in A \times B$ and $\vartheta = \vartheta(\beta)$ is identified.

In order to define super-exogeneity we need to define the class of interventions affecting the DGP. We define the class of interventions under consideration, as any action $\lambda_t \in \Lambda_t$ by an agent from his available action set Λ_t , which alters θ from its current value to a different value $\theta_t = g(\lambda_t, \theta)$. The class of interventions² Λ can then be formally defined as:

$$\Lambda = \{\lambda_t : \theta_t = g(\lambda_t, \theta), \lambda_t \in \Lambda_t\}.$$

We can restrict the class of interventions under consideration by reducing the action set to those describing changes in the marginal process brought about by a policy agency which controls α :

$$\Lambda = \{\lambda_t : \alpha_t = g(\lambda_t, \alpha), \lambda_t \in \Lambda_t\}.$$

DEFINITION 3: *Super-exogeneity: The process z_t is called super exogenous for the parameter of interest $\vartheta(\beta)$ and the class Λ of interventions if there exists a parametrisation of the model such that:*

$$\begin{aligned} f(x_1, \dots, x_T; \alpha_1, \dots, \alpha_T, \beta) \\ = \prod_{t=1}^T g_t(y_t | z_t, x_1, \dots, x_{t-1}; \beta) h_t(z_t | x_1, \dots, x_{t-1}; \alpha_t), \end{aligned}$$

where $(\alpha_1, \dots, \alpha_T, \beta) \in \Lambda \times B$ and $\vartheta = \vartheta(\beta)$ is identified.

2. Among the class of interventions that we will consider are those interventions induced by policy-makers through changes of the level of some variables under their control.

Super-exogeneity establishes conditions under which the Lucas critique can be refuted. Consider y_t as the variable about which agents form plans given all information \mathcal{I}_{t-1} available at time $t - 1$. We would be interested in analyzing the invariance of β when the marginal process changes. FAVERO and HENDRY [1992] distinguishes two levels of the critique applicable to β .

Level A: At this level of the critique, β may change due to changes in economic policy control rules.

Level B: At this level of the critique, β might vary because of changes in economic environment that alter expectations.

At *level A*, changes in the distribution of variables that are under the control of a policymaker lead to variations in the parameters of empirical models. In this case future expectations are not necessarily involved. A case of this critique is the LUCAS [1975] supply function. The *level B* of the critique corresponds to the use of backward-looking econometric specifications when agents use forward-looking specifications.

5.2 The Model

Consider the VAR(p) in Equation (1). If we apply the partition $x_t = (y_t' : z_t)'$ and consider just one lag, $p = 1$, we have:

$$(15) \quad \begin{pmatrix} y_t \\ z_t \end{pmatrix} = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \begin{pmatrix} y_{t-1} \\ z_{t-1} \end{pmatrix} + \begin{pmatrix} \mu_{y,t} \\ \mu_{z,t} \end{pmatrix} + \begin{pmatrix} \varepsilon_{y,t} \\ \varepsilon_{z,t} \end{pmatrix}.$$

If we consider all information as of time $t - 1$ and denote it by \mathcal{I}_{t-1} , the unrestricted model can be written as:

$$(16) \quad \begin{pmatrix} y_t \\ z_t \end{pmatrix} \Big| \mathcal{I}_{t-1} \sim \text{NID} \left(\begin{pmatrix} \mu_{y,t}^c \\ \mu_{z,t}^c \end{pmatrix}, \begin{pmatrix} \Sigma_{yy} & \Sigma_{yz} \\ \Sigma_{zy} & \Sigma_{zz} \end{pmatrix} \right),$$

where $\mu_{y,t}^c := E(y_t | \mathcal{I}_{t-1})$, $\mu_{z,t}^c := E(z_t | \mathcal{I}_{t-1})$ and the intercept μ_t is subject to regime shifts.

HENDRY and MIZON [1998] have advanced two different situations in which common deterministic shifts could play an essential role in modelling. They refer to these situations as the *contemporaneous correlation* case and the *behavioral relation* case. The contemporaneous and behavioral cases can be identified with the *level A* and *level B* of the *Lucas critique* presented above. In the *contemporaneous correlation case* y_t can be seen as a policy variable whereas z_t is an instrument that policymakers can use in order to reach their goal in terms of y_t . The *behavioral relation case* refers to the situation in which agents form rational expectations (about z_t) and there is an interest in analyzing how changes in the expectations may affect the plan of the agents

(y_t) .³ In both cases, common deterministic shifts are introduced to justify invariance of the conditional model due to changes in the marginal model. The existence of a specific linear relationship relating breaks $(\Sigma_{yz} \Sigma_{zz}^{-1})$ under the presence of weak exogeneity (see ENGLE *et al.*, [1983]) define necessary conditions for a valid policy analysis.

In order to illustrate the empirical importance of the *Lucas critique* and the empirical applicability of the super-exogeneity test that we will propose later, we will briefly discuss an economic example of the *level A* of the critique. We mentioned above that a case of *level A* of the critique is the Lucas supply function. In the model presented in LUCAS [1975] and discussed in LUCAS [1976], the economy is considered as formed by suppliers of goods distributed over N distinct markets $i = 1, \dots, N$. The quantity supplied in each market is composed of two components:

$$y_{it} = y_{it}^p + y_{it}^c,$$

where y_{it}^p and y_{it}^c are respectively the permanent and transitory components of output supplied by individual i to the market. Transitory supply varies with the relative prices of goods in market i :

$$(17) \quad y_{it}^c = \beta(p_{it} - p_{it}^e),$$

where p_{it} and p_{it}^e are the local and general level of prices in the economy. The local level of prices in market i (p_{it}) consist of two components:

$$p_{it} = p_t + z_{it}.$$

Suppliers do not observe these components separately and just observe p_{it} . Conditioned on all information prior to time t (\mathcal{I}_{t-1}), p_t is assumed to be normally distributed with mean \bar{p}_t and variance σ^2 :

$$p_t \sim \text{NID}(\bar{p}_t, \sigma^2).$$

The component z_{it} is independent across time and across markets and is distributed normally with mean 0 and variance τ^2 :

$$z_{it} \sim \text{NID}(0, \tau^2).$$

The general price is an average over all markets and considering a law of large number for z_{it} , the general price level is given by p_t . The estimate of the conditional mean of p_t is p_{it}^e and can be obtained as:

$$(18) \quad p_{it}^e = E[p_t | \mathcal{I}_{t-1}, p_{it}] = (1 - \theta)p_{it} + \theta\bar{p}_t \text{ with } \theta = \frac{\tau^2}{\tau^2 + \sigma^2}.$$

3. Many models in economics are expressed in terms of rational expectations. They can be expressed as the behavioral relation:

$$E(y_t | \mathcal{I}_{t-1}) = \mu_y^* + \Psi E(z_t | \mathcal{I}_{t-1})$$

or in short form: $\mu_{y,t} = \mu_y^* + \Psi \mu_{z,t}$, where $\mu_{y,t}$ are the agent's plan about variables they control and $\mu_{z,t}$ are the policymakers plans about which the agents hold rational expectations. Changes in the marginal process will induce changes in the conditional model but it can be shown that the presence of common deterministic shifts could lead to a conditional model with invariant parameters.

Substituting expression for p_{it}^e in (17), averaging over all markets and adding up the permanent component we have the *Lucas* supply function:

$$y_t = \theta\beta(p_t - \bar{p}_t) + y_{pt}.$$

In the Lucas supply function, output is viewed as made up of a permanent component and a transitory component that depends on deviations of the current level of nominal prices from the expected level given \mathcal{I}_{t-1} . The expected price level conditional on past information \bar{p}_t will vary with the average inflation rate.

LUCAS [1976] emphasized that though the econometrician might infer a stable trade-off between (transitory) output and the level of inflation, whenever this trade-off was exploited, the relationship broke down. Assume the price level follows a random walk with inflation a stationary process:

$$p_t = p_{t-1} + \varepsilon_t,$$

with $\varepsilon_t \sim \text{NID}(\pi, \sigma^2)$. Conditional on all information at time $t-1$, $\bar{p}_t = p_{t-1} + \pi$, and the relation between output and inflation is given by:

$$(19) \quad y_t = \theta\beta(p_t - p_{t-1}) - \theta\beta\pi + y_{pt}.$$

If policymakers want to exploit an empirical relationship such as that implied by (19) and alter the rate of inflation π , the relationship that initially seemed characterized by stable parameters will be subject to shifts.

Policy-induced changes in the distribution of the marginal process lead to changes in the conditional model but the invariance of the parameters of interest is guaranteed in the presence of super-exogeneity. In an empirical model of output and inflation, shifts in the marginal model for inflation induced by policymakers, will induce changes in the parameters of the conditional model for output unless super-exogeneity holds. Shifts in the conditional and marginal process occur simultaneously and thus interventions could be modeled within the framework we will discuss in the following. A CDS test for super-exogeneity could be implemented to analyze the invariance of the conditional parameters and will be presented in the next section.

5.3 The Conditional System

Valid inference from a conditional system requires weak exogeneity of the marginal process with respect to the parameters of interest in the conditional model. Using the normality of ε_t , the reduced-form model in (15) can be expressed in terms of the conditional model and the marginal model as:

$$(20) \quad y_t \mid z_t, \mathcal{I}_{t-1} \sim \text{NID}\left(\mu_{y|z,t}^c, \Omega\right),$$

$$(21) \quad z_t \mid \mathcal{I}_{t-1} \sim \text{NID}\left(\mu_{z,t}^c, \Sigma_{zz}\right),$$

where the density of y_t conditional on z_t and \mathcal{I}_{t-1} is determined by:

$$\begin{aligned}\mu_{y|z,t}^c &: = \mathbb{E}(y_t | z_t, \mathcal{I}_{t-1}) = \mu_{y,t} + \Sigma_{yz} \Sigma_{zz}^{-1} (z_t - \mu_{z,t}), \\ \Omega &: = \text{Var}(y_t | z_t, \mathcal{I}_{t-1}) = \Sigma_{yy} - \Sigma_{yz} \Sigma_{zz}^{-1} \Sigma_{zy}.\end{aligned}$$

Rewriting (15) in terms of the conditional and marginal model results in:

$$(22) \quad \begin{pmatrix} \mathbf{I} & -\Sigma_{yz} \Sigma_{zz}^{-1} \\ \mathbf{0} & \mathbf{I} \end{pmatrix} \begin{pmatrix} y_t \\ z_t \end{pmatrix} = \begin{pmatrix} \mathbf{I} & -\Sigma_{yz} \Sigma_{zz}^{-1} \\ \mathbf{0} & \mathbf{I} \end{pmatrix} \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \begin{pmatrix} y_{t-1} \\ z_{t-1} \end{pmatrix} \\ + \begin{pmatrix} \mathbf{I} & -\Sigma_{yz} \Sigma_{zz}^{-1} \\ \mathbf{0} & \mathbf{I} \end{pmatrix} \begin{pmatrix} \mu_{yt} \\ \mu_{zt} \end{pmatrix} + \begin{pmatrix} u_{y,t} \\ u_{z,t} \end{pmatrix},$$

where the variance matrix of the transformed residuals is block diagonal:

$$u_t \sim \text{NID} \left(\begin{pmatrix} \mathbf{0} \\ \mathbf{0} \end{pmatrix}, \begin{pmatrix} \Omega & \mathbf{0} \\ \mathbf{0} & \Sigma_{zz} \end{pmatrix} \right).$$

In general, this type of model is prone to suffer from the *Lucas critique*. That is, changes in the marginal model lead to non-constancy of the conditional model. Shifts of the marginal model may induce shifts in the conditional model, but a convenient linear combination can induce constancy in the conditional process, such that:

$$(23) \quad (\mathbf{I} : -\Sigma_{yz} \Sigma_{zz}^{-1}) \begin{pmatrix} \mu_{yt} \\ \mu_{zt} \end{pmatrix} = \text{constant}.$$

If we model $(\mu'_{yt} : \mu'_{zt})'$ with the corresponding intervention variables, Equation (15) can be rewritten as:

$$(24) \quad x_t = A_1 x_{t-1} + \mathcal{M} D_t + \varepsilon_t,$$

Note that the condition in (23) requires a linear relationship of breaks across equations. Hence, in the presence of super-exogeneity, we would have a reduced rank condition on the coefficients of the intervention variables used to model the unrestricted system (\mathcal{M}), such that we can rewrite the previous model as:

$$(25) \quad x_t = A_1 x_{t-1} + \eta \xi' D_t + \varepsilon_t.$$

Furthermore, in order for super-exogeneity to hold, the conditional model should be invariant to the set of interventions in the marginal process, which would require:

$$(26) \quad (\mathbf{I} : -\Sigma_{yz} \Sigma_{zz}^{-1}) \eta = 0.$$

We could define $C = (\mathbf{I} : -\Sigma_{yz} \Sigma_{zz}^{-1})$ as the $n_y \times (n_y + n_z)$ matrix that allows us to recover the conditional model from the full model, where n_y and n_z define the dimension of the vector y_t and z_t respectively. Given that η is a $(n_y + n_z \times r)$ matrix, we then have that η_{\perp} is a $(n_y + n_z \times n_y + n_z - r)$

matrix. In order for $C \subset \eta'_\perp$ and $C\eta = 0$, one needs $n_y + n_z - r \geq n_y$ or $n_z \geq r$.⁴ So the reduced rank of the coefficient of the intervention dummies with specific restrictions on the null space of η and weak exogeneity imply super-exogeneity. This postulates the invariance of the conditional model under a set of interventions in the marginal model.

6 Testing for Super-Exogeneity

The previous subsection showed how super-exogeneity of the y_t process with respect to a set of interventions (shifts in the conditional mean of the marginal process) required a reduced rank condition of the coefficients of the intervention variables. In order to implement a likelihood ratio test for super-exogeneity with respect to this class of interventions, we firstly need to estimate the model under the null characterized by a reduced rank of $\mathcal{M} = \eta\xi'$ and specific restrictions on C , $C \subset \eta'_\perp$; and then under the alternative, *ie.* the unrestricted model with the reduced rank of $\mathcal{M} = \eta\xi'$ imposed. In what follows, we present the original procedure to test for super-exogeneity proposed by ENGLE and HENDRY [1993] and two alternative new procedures. These three procedures differ in the way in which the model is estimated under the null and how the test is constructed. The first procedure reparameterizes the original approach of ENGLE and HENDRY [1993] as a reduced rank restriction where the null corresponds to ENGLE and HENDRY [1993], but the alternative is defined differently. The second method involves reduced-rank-regression estimations of the reduced-form model under the conditions of super-exogeneity and $n_z = r$. The third method introduces a very general approach to the restricted estimation of η and ξ , where additional restrictions can be imposed.

6.1 The Engle and Hendry Procedure

A simple testing procedure can be implemented with just linear regressions. In order to show this procedure let us start from the model in (15):

$$(27) \quad \begin{pmatrix} y_t \\ z_t \end{pmatrix} = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \begin{pmatrix} y_{t-1} \\ z_{t-1} \end{pmatrix} + \mathcal{M} \begin{pmatrix} d_{t1} \\ d_{t2} \end{pmatrix} + \begin{pmatrix} \varepsilon_{y,t} \\ \varepsilon_{z,t} \end{pmatrix}.$$

where \mathcal{M} is a matrix of coefficients of the intervention variables,

$$\mathcal{M} = \begin{pmatrix} \mu_1^y & \mu_2^y \\ \mu_1^z & \mu_2^z \end{pmatrix}.$$

4. When y_t and z_t do not make up the full model the rank of η could be greater than n_z . We can exemplify this using a trivariate system, where $x_t = (y_t, z_t, w_t)'$ and the $\text{rank}(\eta) = 2$. In this case one might want to investigate the super-exogeneity of z_t with respect to a class of interventions Λ in the conditional model $y_t|z_t$. We could then test the hypothesis $H_0 : \eta = (\eta_1 : \eta_2)$, where η_1 and η_2 are 3×1 vectors with $\eta_1 = (\sigma_{zz}^{-1}\sigma_{zy}, 1, 0)'$ and η_2 can be left unrestricted. Testing this type of restrictions is discussed in Subsection 6.3.

The conditional model is given by:

$$\begin{aligned} y_t &= A_{11}y_{t-1} + A_{12}z_{t-1} + \mu_1^y d_{t1} + \mu_2^y d_{t2} + E(\varepsilon_{y,t} | \varepsilon_{z,t}), \\ &= A_{11}y_{t-1} + A_{12}z_{t-1} + \mu_1^y d_{t1} + \mu_2^y d_{t2} \\ &\quad + \omega(z_t - A_{21}y_{t-1} - A_{22}z_{t-1} - \mu_1^z d_{t1} - \mu_2^z d_{t2}) + \tilde{\varepsilon}_{y,t}, \end{aligned}$$

with $\omega = \Sigma_{yz} \Sigma_{zz}^{-1}$ and $\tilde{\varepsilon}_{y,t} = \varepsilon_{y,t} - \omega \varepsilon_{z,t}$, which can be rewritten as:

$$\begin{aligned} y_t &= \omega z_t + (A_{11} - \omega A_{21})y_{t-1} + (A_{12} - \omega A_{22})z_{t-1} + (\mu_1^y - \omega \mu_1^z) d_{t1} \\ &\quad + (\mu_2^y - \omega \mu_2^z) d_{t2} + \tilde{\varepsilon}_{y,t}, \end{aligned}$$

and the marginal model is given by:

$$(28) \quad z_t = A_{21}y_{t-1} + A_{22}z_{t-1} + \mu_1^z d_{t1} + \mu_2^z d_{t2} + \varepsilon_{z,t}.$$

Under the super-exogeneity condition, we have the restrictions $(\mu_1^y - \omega \mu_1^z) = \mathbf{0}$ and $(\mu_2^y - \omega \mu_2^z) = \mathbf{0}$, which are the reduced rank conditions of Equation (23). For bivariate models with the restriction that $\text{rank}(\mathcal{M}) = n_z = 1$, these restrictions can also be written as $\mu_1^y / \mu_1^z = \mu_2^y / \mu_2^z = \omega$, where the specific restrictions on the null space of η become clearer. Under the null of super-exogeneity, we have that the conditional process reduces to

$$(29) \quad y_t = \omega z_t + (A_{11} - \omega A_{21})y_{t-1} + (A_{12} - \omega A_{22})z_{t-1} + \tilde{\varepsilon}_{y,t}.$$

The parameters in the conditional model are $\theta_c = \{\omega, A_{11} - \omega A_{21}, A_{12} - \omega A_{22}, \Sigma_{yy} - \Sigma_{yz} \Sigma_{zz}^{-1} \Sigma_{zy}\}$ and the parameters in the marginal model are $\theta_m = \{A_{21}, A_{22}, \mu_1^z, \Sigma_{zz}\}$. The Gaussianity of the errors implies that the parameters in the marginal process are variation free of the parameters in the conditional process. Equations (29) and (28) can be estimated separately and the full maximum likelihood estimate is made up of two factors corresponding to the marginal and conditional density. The maximum likelihood estimation under the alternative just requires the estimation of the model under the reduced rank restriction along the lines of Section 3.⁵ The construction of a likelihood ratio test is thus straightforward.

6.2 The CDS Testing Procedure

In the special case in which $n_z = r$ we could parametrize η as $\eta' = (\Sigma_{zz}^{-1} \Sigma_{zy} : \mathbf{I})'$.⁶ In order to show the estimation procedure of the model under these conditions let us start from the model equation:

$$R_X = \mathcal{M} R_D + E,$$

5. This estimation procedure could be used when $n_z > r$.

6. Note that $\eta = (\Sigma_{zz}^{-1} \Sigma_{zy} : \mathbf{I})'$ is not the only possible parametrisation and for any $r \times r$ full rank matrix, say B , we could also express η as $\eta = (\Sigma_{zz}^{-1} \Sigma_{zy} : \mathbf{I})' B$.

where R_X and R_D are the corrected residuals from Equation (4). Imposing the reduced rank restriction $\mathcal{M} = \eta\xi'$ we get,

$$(30) \quad R_X = \eta\xi' R_D + E.$$

The maximum likelihood of this model under the super-exogeneity restriction can be calculated as follows:

1. Start with initial estimates $\{\eta^{(0)}, \Sigma^{(0)}\}$ of η and Σ satisfying the restriction:

$$\eta^{(0)'} = \left(\Sigma_{zz}^{(0)-1} \Sigma_{zy}^{(0)} : \mathbf{I} \right).$$

2. Multiplying (30) through by $\bar{\eta}^{(0)'} = \left(\eta^{(0)'} \eta^{(0)} \right)^{-1} \eta^{(0)'}$ results in an ordinary linear regression problem in ξ :

$$\bar{\eta}^{(0)'} R_X = \xi' R_D + \bar{\eta}^{(0)'} E.$$

Applying OLS to the previous equation delivers new estimates of ξ , denoted $\xi^{(1)}$.

3. Given $\{\xi^{(1)}, \Sigma^{(0)}\}$, new estimates of η and Σ can be obtained from (30):

$$R_X = \eta \left(\xi^{(1)'} R_D \right) + E.$$

We can loop in this algorithm till we obtain convergence, where in each iteration the restriction $\eta' = \left(\Sigma_{zz}^{-1} \Sigma_{zy} : \mathbf{I} \right)$ is always updated.

6.3 A Generalized CDS Testing Procedure

An alternative estimation procedure of the model under the super-exogeneity restriction can be based on the first-order conditions of the maximum likelihood problem. Let us start from the concentrated likelihood function,

$$\begin{aligned} \ln L = & -\frac{nT}{2} \ln 2\pi - \frac{T}{2} \ln |\Sigma| \\ & - \frac{1}{2} \text{tr} \Sigma^{-1} \left[(S_{XX} - S_{XD} \xi \eta' - \eta \xi' S_{DX} + \eta \xi' S_{DD} \xi \eta') \right]. \end{aligned}$$

Introducing a very general formulation, we can write the linear restriction for ξ and η as:

$$\text{vec}(\xi) = H\varphi + h,$$

$$\text{vec}(\eta') = G\chi.$$

Where ξ denotes the matrix of the linear combinations of the intervention variables with identifying restrictions imposed. H is a known $nr \times k$ matrix and $\varphi \in R^k$ is an unrestricted parameter vector. The vector h corresponds to

normalization restrictions on the vector of ξ . G is a known $nr \times l$ matrix of full rank and $\chi \in R^l$ is an unrestricted parameter vector. The purpose of this formulation is that we can impose jointly further restrictions to those implied by the super-exogeneity restrictions. The two previous procedures dealt with the bivariate case. Now, we exemplify this procedure using a trivariate system, where $x_t = (y_t, z_t, w_t)'$. One might want to investigate the super-exogeneity of w_t with respect to a class of interventions Λ in the conditional model $y_t|w_t$. We have seen in the previous section that this implies a set of restrictions on η . With this procedure one could test the super-exogeneity restrictions jointly with further restrictions on the coefficients of the loading (η) on z_t or restrictions on ξ .

The derivatives of the likelihood function with respect to φ and χ are given by:

$$(31) \quad \frac{\partial \ln L}{\partial \varphi} = H' \text{vec}(S_{DX} \Sigma^{-1} \alpha) - H' \text{vec}(S_{DD} \xi \eta' \Sigma^{-1} \eta).$$

$$(32) \quad \frac{\partial \ln L}{\partial \chi} = G' \text{vec}(\xi' S_{DX} \Sigma^{-1}) - G' \text{vec}(\xi' S_{DD} \xi \eta' \Sigma^{-1}).$$

By substituting the restrictions of ξ in (31) we get:

$$(33) \quad \varphi(\chi, \Sigma) = \left[H'(\eta' \Sigma^{-1} \eta \otimes S_{DD}) H \right]^{-1} \left[H'(\eta' \Sigma^{-1} \otimes \mathbf{I}) \text{vec}(S_{DX}) - H'(\eta' \Sigma^{-1} \eta \otimes S_{DD}) h \right].$$

Similarly for χ , we can substitute the restrictions for η in (32) and we get:

$$(34) \quad \chi(\varphi, \Sigma) = \left[G'(\Sigma^{-1} \otimes \xi' S_{DD} \xi) G \right]^{-1} \left[G'(\Sigma^{-1} \otimes \xi') \text{vec}(S_{DX}) \right].$$

It can easily be seen that the first-order condition for Σ , for given χ and φ , is given by:

$$(35) \quad \Sigma(\varphi, \chi) = S_{XX} - S_{XD} \xi \eta' - \eta \xi' S_{DX} + \eta \xi' S_{DD} \xi \eta'.$$

Hence, for initial χ and Σ we can impose the restrictions on η and obtain estimates of ξ from (33). For given ξ and Σ new estimates of χ can be obtained from (34). For given ξ and η , equation (35) delivers new estimates of Σ . We can then iterate in this algorithm with the corresponding super-exogeneity restrictions always updated (see footnote 4).

The likelihood ratio test can be shown to have a χ^2 distribution. The degrees of freedom result from comparing the tangent space of $\eta \xi'$ with and without restrictions (see JOHANSEN [1995]).

7 A Small Monte Carlo Experiment

The power of different tests for super-exogeneity has been analyzed by PSARADAKIS and SOLA [1996] and FAVERO and HENDRY [1992] for *level B* of the *Lucas critique*: while agents are forward-looking, the econometric model is based on backward-looking expectations, and changes in the economic environment alter expectations inducing changes in the conditional model. The power results obtained in their simulation experiment lead PSARADAKIS and SOLA [1996] to question the usefulness of super-exogeneity and invariance tests for the assessment of the empirical relevance of forward-looking models. While FAVERO and HENDRY [1992] also question the usefulness of the tests for the assessment of forward-looking models, they interpret the Monte Carlo results as a lack of strength of the *Lucas critique*: “for changes of a magnitude that are large enough to be policy relevant, conditional models approximating expectations process do not in practice experience dramatic predictive failure” (p. 290). In contrast to the previous literature, we focus on the power of super-exogeneity test at *level A* of the *Lucas critique*.⁷ The following Monte Carlo study is designed to analyze the size and power of the CDS tests for super-exogeneity proposed in Section 6.2. We will compare the results with the test introduced by ENGLE and HENDRY [1993].

In the Monte Carlo simulations, we use the benchmark DGP_2 of Section 4 with a reduced-rank matrix of intervention coefficients:

$$(36) \quad \begin{pmatrix} y_t \\ z_t \end{pmatrix} = \begin{pmatrix} 0.75 & 0.5 \\ 0 & 0.8 \end{pmatrix} \begin{pmatrix} y_{t-1} \\ z_{t-1} \end{pmatrix} + \begin{pmatrix} \mu_1^y & \mu_2^y \\ \mu_1^z & \mu_2^z \end{pmatrix} \begin{pmatrix} d_{t1} \\ d_{t2} \end{pmatrix} + \begin{pmatrix} \varepsilon_{1t} \\ \varepsilon_{2t} \end{pmatrix}$$

where $\begin{pmatrix} \varepsilon_{1t} \\ \varepsilon_{2t} \end{pmatrix} \sim \text{NID} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \sigma_{yy} & \sigma_{yz} \\ \sigma_{zy} & \sigma_{zz} \end{pmatrix} \right)$.

The break points are again at point $\tau_1 = 0.3$ and $\tau_1 = 0.7$ of the sample. Under super-exogeneity the restrictions that we will impose in the matrix \mathcal{M} will imply restrictions in the matrix Ω , and this resulting matrix will be our new Ω in the benchmark DGP_2 .

We will first consider the case in which the shifts in the marginal process z_t also affect the y_t process, but the parameters of the conditional process $y_t|z_t$ remain constant. The corresponding restrictions on the intervention coefficients are given by the reduced rank restriction, $\mathcal{M} = \eta\xi'$, and the super-exogeneity restriction $\eta' = (\Sigma_{zz}^{-1}\Sigma_{zy} : \mathbf{I})$. For our bivariate case, these restrictions can be written as:

$$(37) \quad \frac{\mu_1^y}{\mu_1^z} = \frac{\mu_2^y}{\mu_2^z} = \frac{\sigma_{zy}}{\sigma_{zz}}.$$

7. It is worth mentioning that in most settings when expectations are modeled as forward-looking, weak exogeneity of the marginal process with respect to the parameters of interest is generally violated. In this case, testing for the constancy of the parameters of the conditional process or additional variable tests for the absence of the moments of the marginal process in the conditional models are invariance tests.

Under condition (37), the parameters in the conditional process are invariant to the class of interventions in the marginal process. In other words, despite changes in the marginal distribution induced by the policymaker's alteration of his policy rule (here, intercept shift), the parameters in the conditional model are invariant to this intervention. The *Lucas critique* does not hold for this specific parametrisation of the DGP. Let us call this model DGP_1^* .

Alternatively, we consider the case where the parameters of the conditional model are not invariant to changes in the intercept of the policy rule. The intervention matrix $\mathcal{M} = \eta \xi'$ still has reduced rank, but the super-exogeneity condition does not hold:

$$\frac{\mu_1^y}{\mu_1^z} = \frac{\mu_2^y}{\mu_2^z} = \frac{\sigma_{zy}}{\sigma_{zz}} + h,$$

with $h \neq 0$. In the simulations we will use a value of 0.5 for h . This correspond to a situation in which shifts in the y_t and z_t process are linearly related, the matrix of the intervention coefficients is of reduced rank but super-exogeneity does not hold. So the conditional model has non-constant parameters. We will refer to this case as DGP_2^* .

In the previous section, it has been shown that the test for super-exogeneity is a test of $H_0: \text{rank}(\mathcal{M}) = 1 \wedge \eta' = (\sigma_{zz}^{-1} \sigma_{zy}: 1)$ against $H_1: \text{rank}(\mathcal{M}) = 1 \wedge \eta' \neq (\sigma_{zz}^{-1} \sigma_{zy}: 1)$, which follows a $\chi^2(1)$. In order to analyze the size of the test, we will consider DGP_1^* for sample sizes of $T = 50, 100$ and 150 .

Size and power of the CDS test for super-exogeneity are reported in Tables 3 and 4. Table 3 presents the rejection frequencies for the CDS test for super-exogeneity of $H_0: \text{rank}(\mathcal{M}) = 1 \wedge \eta' = (\sigma_{zz}^{-1} \sigma_{zy}: 1)$ against $H_1: \text{rank}(\mathcal{M}) = 1 \wedge \eta' \neq (\sigma_{zz}^{-1} \sigma_{zy}: 1)$ under the null (DGP_1^*). Table 4 shows the rejection frequencies of the test when the simulated model is DGP_2^* . The actual size of the CDS test for super-exogeneity differs only slightly from nominal size and power (non size corrected) is high.

For comparison purposes, we also consider the ENGLE and HENDRY [1993] test for super-exogeneity. This is a variable addition tests constructed as a conventional F -test for the joint significance of the intervention variables in the conditional model: $H_0: \mu_{1,c}^y = \mu_{2,c}^y = 0$ against $H_1: \mu_{1,c}^y \neq 0$ and $\mu_{2,c}^y \neq 0$.⁸ The results for size and power of this test are presented in Tables 5 and 6. Table 5 reports the rejection frequencies for the test on the conditional model of $H_0: \mu_{1,c}^y = \mu_{2,c}^y = 0$ against $H_1: \mu_{1,c}^y \neq 0$ and $\mu_{2,c}^y \neq 0$ when the simulated model is DGP_1^* . Table 6 presents the rejection frequencies of the test when the simulated model is DGP_2^* . The size and (non-size corrected) power results obtained for the Engle and Hendry test for super-exogeneity are similar to the ones for the CDS test, though there is some indication that the *Engle* and *Hendry* test does not perform as good as the CDS test.

8. Where $\mu_{1,c}^y = (\mu_1^y - \omega \mu_1^z)$ and $\mu_{2,c}^y = (\mu_2^y - \omega \mu_2^z)$ are the coefficients of the intervention variables in the conditional model.

TABLE 3
Size of the CDS Test for Super-Exogeneity

T	20%	10%	5%	1%
50	0.298	0.183	0.114	0.0377
100	0.266	0.153	0.086	0.0232
150	0.256	0.147	0.081	0.0217

Note: Rejection frequencies for the CDS test statistic for DGP_1^* .

TABLE 4
Power of the CDS Test for Super-Exogeneity

T	20%	10%	5%	1%
50	0.971	0.948	0.917	0.809
100	0.997	0.994	0.988	0.958
150	0.999	0.999	0.997	0.989

Note: Rejection frequencies for the CDS test statistic for DGP_2^* .

TABLE 5
Size of the Engle and Hendry Test for Super-Exogeneity

T	20%	10%	5%	1%
50	0.349	0.216	0.129	0.0353
100	0.331	0.202	0.121	0.0323
150	0.309	0.184	0.106	0.0282


Note: Rejection frequencies for the Engle and Hendry test for DGP_1^* .

TABLE 6
Power of the Engle and Hendry Test for Super-Exogeneity

T	20%	10%	5%	1%
50	0.971	0.943	0.901	0.753
100	0.997	0.992	0.983	0.941
150	0.999	0.998	0.996	0.982

Note: Rejection frequencies for the Engle and Hendry test for DGP_2^* .

FIGURE 6
Size-Power Curve Analysis for the Engle-Hendry and CDS Tests for Super-Exogeneity



In order to be able to compare the properties of the two tests further, size-power curves are plotted in Figure 6 for both tests for sample sizes of $T = 50$ and 100. The left hand panel of Figure 6 shows the results for a sample size of 50. The continuous line graph corresponds to the size-power curve for the CDS and the discontinuous line corresponds to the *Engle* and *Hendry* test. These graphs show that for a given size the CDS test for super-exogeneity has a higher power than the *Engle* and *Hendry* test. For a sample size of 100 (RHS of Figure 6), the behavior of the CDS test is still better, though the differences between the behavior are greatly reduced. For larger samples, results did not differ much.

In the light of these results, the use of the CDS tests for super-exogeneity in empirical applications can be recommended. As we have seen, the CDS test for super-exogeneity has some comparative advantages versus tests proposed in the literature when sample sizes are small, which is often the case in economics.

8 Some Comments

8.1 Empirical Illustrations

In this paper, we propose a general approach to the modelling and testing of common deterministic shifts. An important advantage of this approach is that it does not rely on *prior* information regarding the relationships between the shifts in the mean of the individual processes. This can be useful in cases with limited information about the classification of targets and instruments and their relationship.

Consider, for example, a monetary model where some short-term interest rates (say, the discount rate and the inter-bank rate) are included together with the target variables the policymaker intends to control. The identification of a linear relationship linking shifts in the mean of the different processes would provide valuable information about the exact relationship between instruments and targets, particularly the transmission of interventions (by the weights of the linear relationship that govern the deterministic shifts).

It is worth emphasizing that even in the absence of super-exogeneity, the presence of CDS will deliver important information about the economic system and testing for CDS could be used to ask relevant economic questions. Let us take as an example the defense of a currency by the government. Suppose that the researcher uses a vector autoregression to model the exchange rate, the yield spread of domestic government bonds with a benchmark risk-free bond, and the short-term interest rate as the policy instrument. In a period of exchange rate turbulence, he would probably find that the government has intervened repeatedly raising the interest rate in order to defend the currency. Furthermore suppose that there have been just three interventions and that these actions of the government are represented by intervention dummies. The researcher could then hypothesize that if each of

the interventions was credible, *ceteris paribus*, they would have a proportional effect on the exchange rate and the yield spreads. This would imply a reduced rank of the matrix of dummy coefficients. In contrast, a full rank would indicate the non-credibility of the government interventions (the first intervention had a proportionally different effect than the second and third intervention).

Another example from industrial organization is the analysis presented in GUERRERO, PEÑA and PONCELA [1998]. GUERRERO *et al.* [1998] aim to measure the effects of promotional campaigns launched by a bank in order to attract deposits. The banking institution wanted to identify the possible effects on (a) new accounts, (b) stock variation, (c) cancellations, and (d) total amount. Since the increase in total amount from the previous month to the current month is given by new accounts + stock variation – Cancellation, a linear restriction is satisfied by the vector of time series every month and also applies to the intervention coefficients.

8.2 Limitations, Extensions and Generalizations

The methodology developed in this paper is restricted by two major assumptions. First that, conditional on the breaks, the system is stationary which excludes integrated-cointegrated systems. Secondly, that the breaks points are known *a priori*.

In the case of cointegration, one could be interested in the following model:

$$(38) \quad \Delta x_t = \Pi x_{t-1} + \mathcal{M} D_t + \varepsilon_t,$$

where x_t is $I(1)$, Π has reduced rank and \mathcal{M} and D_t had been defined in section 2. The above model equation raises two different issues.

- The first issue is the mapping of the $I(1)$ system to the $I(0)$ system.
- The second issue is the test of CDS in the vector equilibrium representation of the system. That is, when we have mapped the system from $I(1)$ to $I(0)$.

The second issue is probably the easiest of both issues and we will address it first. A general result is that when the $I(1)$ system has been mapped to $I(0)$, a test of restrictions on η have a χ^2 distribution. In most empirical applications economic priors are used to map the system to $I(0)$ (see FAVERO and HENDRY [1992] for a situation in which an $I(1)$ system is mapped to an $I(0)$ system and then the *Lucas critique* is investigated).

If we map the $I(1)$ system into the vector equilibrium representation:

$$(39) \quad \Delta x_t = \alpha \beta' x_{t-1} + \eta \xi' D_t + \varepsilon_t,$$

the system can then be estimated by a switching algorithm which provides reduced-rank estimations of (i) α, β conditional on η, ξ and (ii) η, ξ conditional on α, β . Thus one would combine the techniques developed in this paper with the well-established cointegration analysis of JOHANSEN [1995] (see TORO [1999]). So, for known rank of Π and \mathcal{M} , one could impose a CDS restriction together with restrictions in α or β and these are χ^2 tests.

A probably more technical and complicated issue is the mapping of the

$I(1)$ to the $I(0)$ system. In order to do this we need the joint determination of the rank on Π and \mathcal{M} . This issue is quite involved. We will try to summarize the main issues without introducing new notation and going into much technicality. Here one could investigate the determination of the rank of Π in the presence of intervention variables and then investigate the rank of \mathcal{M} for given $rank(\Pi) = r$. We first review the literature on the determination of the rank of Π in the presence of intervention variables and then discuss the difficulties involved when \mathcal{M} has also reduced rank suggesting potential solutions to this problem in empirical applications.

Most empirical applications that use the cointegrated VAR system with intervention variables assess the rank of Π based on the traditional tables tabulated in JOHANSEN [1995]. However JOHANSEN and NIELSEN [1993] showed that in the presence of deterministic dummy variables in the cointegrated VAR system, different distributions apply. These are generalized Dickey-Fuller distributions. JOHANSEN and NIELSEN [1993] propose a program for the simulation of the specific distribution. JOHANSEN, NIELSEN and MOSCONI [2000] analyze cointegration in Gaussian vector autoregressive model with a broken linear trend and known break points. They investigate three models which they denote by H_l , H_{lc} and H_c . In model H_l in each subsample the deterministic component is linear both for non-stationary and cointegrating relationships. Model H_c has no linear trend but a broken constant level. In model H_{lc} the non-stationary relations have a broken linear trend whereas the cointegration relations have broken constant levels. They demonstrate that in model H_{lc} , the third model, the asymptotic analysis is heavily burdened with nuisance parameters. The asymptotic distribution is investigated for each model and the moments of the distributions are summarized with a response surface. They suggest to approximate the critical values with a Γ -distribution using the moments obtained in a response surface analysis.

A further possible model is a cointegrated VAR model where the set of intervention coefficients could also have a reduced rank structure. In this model one would be interested in investigating the joint determination of the rank of Π and \mathcal{M} . It has been shown (see TORO [1999]) that the asymptotic distribution of the test for the rank of Π when the regressors of \mathcal{M} are deterministic depends on the rank of \mathcal{M} or the number of deterministic trends that load into the system and whether $\alpha_{\perp}\mathcal{M} = 0$ or $\alpha_{\perp}\mathcal{M} \neq 0$. Also tests on the rank of \mathcal{M} depend on the rank of Π . This seems to be a complicated setting. However, it turns out that a much simpler path exist if the interpretation of the deterministic component is not questioned with the rank but is tested in a later stage (see NIELSEN and RAHBEK [2000]). We could consider the situation in which the intervention variables induce a broken constant level in the cointegrating relationships but they do not induce any broken trend. This is probably the most relevant situation in macroeconomics and it would be a relevant situation in the case of a test of the *Lucas critique*. We would be considering interventions that alter the constant term in an equilibrium relationship but that does not induce a changing trend. This can be formulated as $\alpha'_{\perp}\mathcal{M} = 0$, such that Model Equation 38 can be reparameterized as:

$$(40) \quad \Delta x_t = \alpha(\beta'x_{t-1} + \mathcal{M}_r D_{t-1}) + \mathcal{M}\Delta D_t + \varepsilon_t,$$

with $\mathcal{M}_r = (\alpha' \alpha)^{-1} \alpha' \eta \xi'$. Under the further restriction $\alpha_{\perp} \mathcal{M} = 0$ there is no nuisance parameter in the distribution of the test for the cointegrating rank (see JOHANSEN *et al.* [2000]). Once the rank of Π has been investigated, one could then investigate hypotheses about the rank of $\alpha \mathcal{M}_r = \Upsilon$ or \mathcal{M} in the form of CDS tests or CDS-super-exogeneity test, and they follow a χ^2 distribution.

A further drawback of the reduced rank regression techniques proposed in this paper comes from the fact that the shift points are assumed to be known in advance. If the break points are unknown, then the *Markov*-switching approach provides a powerful tool to model the system.

$$x_t = Ax_{t-1} + \mathcal{M}D_t + \varepsilon_t,$$

where D_t is now a *Markov* chain. KROLZIG [1997] considers the statistical analysis of this system when the (potentially reduced) rank of $\mathcal{M} = \eta \xi'$ is imposed to the system. In an LIML approach, each equation could be estimated separately. Assume that the number of regime is $M = 2$. Then $M - 1$ smoothed regime probabilities associated with each equation can be collected to the matrix D_t and be used in the reduced rank regression approach discussed above. Again one might consider the potential cointegration of the variables, which suggests the combination of including the switching algorithm on each M-step of the EM algorithm for MS-VAR models.

9 Conclusions

This paper brings together two topics of econometric research: common deterministic shifts and super-exogeneity. We have shown how common deterministic shifts can be analyzed with simple and widely-known technique: reduced rank regressions. The proposed CDS tests were found of being only slightly oversized and having excellent power properties in small samples.

Deterministic shifts in the conditional mean of economic variables are a recurrent feature in empirical economics. These shifts happen to affect not just one single economic variable but also contemporaneously, other related variables. Furthermore, these shifts might be related linearly and this linear relationship might prevail throughout time. We have proposed a technique that can be used to analyze such phenomena, and can help to gather important information about how breaks are related through economic variables and across time. Frequently, deterministic shifts are induced by policy changes. Policymakers move the level of some variables in order to affect some target variables and reach specific goals. When deterministic shifts are induced by policymakers, the relationship between common deterministic shifts and super-exogeneity becomes apparent. ▼

• References

- ANDERSON T.W. (1958). – *An Introduction to Multivariate Statistical Analysis*, New York, John Wiley.
- DAVIDSON R., MACKINNON, J. (1997). – « Graphical Methods for Investigating the Size and Power of Hypothesis Tests », *Discussion Paper*.
- ENGLER R.F., GRANGER C.W.J. (1987). – « Co-integration and Error Correction: Representation, Estimation and Testing », *Econometrica*, 55, p. 251-276.
- ENGLER R.F., HENDRY D.F. (1993). – « Testing Super-Exogeneity and Invariance in Regression Models », *Journal of Econometrics*, 56, p. 119-139.
- ENGLER R.F., HENDRY D.F., RICHARD J. (1983). – « Exogeneity », *Econometrica*, 51, p. 277-304.
- ENGLER R.F., KOZICKI S. (1993). – « Testing for Common Features », *Journal of Business Economic and Statistics*, 11, p. 369-395.
- FAVERO C., HENDRY D.F. (1992). – « Testing the Lucas Critique: A Review », *Econometric Reviews*, 11, p. 265-306.
- GUERRERO V., PEÑA D., PONCELA P. (1998). – « Measuring Intervention Effects on Multiple Time Series Subjected to Linear Restrictions: A Banking Example », *Journal of Business Economics and Statistics*, 16, p. 489-497.
- HENDRY D.F. (1997). – « A Theory of Co-Breaking », *Manuscript*, Institute of Economics and Statistics, Oxford.
- HENDRY D.F., MIZON G. (1998). – « Exogeneity, Causality and Co-Breaking in Economic Policy Analysis of a Small Econometric Model of Money in the UK », *Empirical Economics*, 23, p. 267-295.
- JOHANSEN S. (1995). – *Likelihood-Based Inference in Cointegrated Vector Autoregressive Models*, Oxford, Oxford University Press.
- JOHANSEN S., NIELSEN B. (1993). – « Asymptotics for Cointegration Rank Tests in the Presence of Intervention Dummies, Manual for the Simulation Program Disco », *Preprint*, University of Copenhagen, <http://www.math.ku.dk>.
- JOHANSEN S., NIELSEN B., MOSCONI R. (2000). – « Cointegration Analysis in the Presence of Structural Breaks in the Deterministic Trend », *Econometrics Journal*, 3, p. 216-249.
- KROLZIG H.M. (1997). – *Markov Switching Vector Autoregressions, Modelling, Statistical Inference and Application to Business Cycle Analysis*, Berlin: Springer.
- LUCAS R. (1975). – « Some International Evidence on Output-Inflation Tradeoffs », *American Economic Review*, 63, p. 326-334.
- LUCAS R. (1976). – « Econometric Policy Evaluation: A Critique », *Carnegie-Rochester conference series on public policy*, 1, p. 19-46.
- LÜTKEPOHL H. (1991). – *Introduction to Multiple Time Series Analysis*, Berlin: Springer.
- NIELSEN B., RAHBEK A. (2000). – « Similarity Issues in Cointegration Models », *Oxford Bulletin of Economics and Statistics*, 6, p. 5-22.
- PSARADAKIS Z., SOLA M. (1996). – « On the Power of Tests for Superexogeneity and Structural Invariance », *Journal of Econometrics*, 72, p. 151-175.
- TORO J. (1999). – « Modelling Common Deterministic Shifts in $I(1)$ Variables », *Mimeo*, Department of Economics, European University Institute.
- TSO K.S. (1981). – « Reduced-Rank Regression and Canonical Analysis », *Journal of the Royal Statistical Society B*, 43, p. 183-189.

